



Discover Generics

Cost-Effective CT & MRI Contrast Agents



FRESENIUS
KABI

[VIEW CATALOG](#)

AJNR

Congenital malignant gliosarcoma.

S J Goldstein, B Young and W R Markesberry

AJNR Am J Neuroradiol 1981, 2 (5) 475-476

<http://www.ajnr.org/content/2/5/475.citation>

This information is current as
of September 5, 2025.

Congenital Malignant Gliosarcoma

Steven J. Goldstein,¹ Byron Young,² and William R. Markesberry³

Malignant gliosarcoma is an extremely rare, anaplastic tumor which to date has been found exclusively in adults. This type of sarcoma is a form of mixed glioma and sarcoma which is quite different and distinct from the type of sarcoma that arises in a glioblastoma [1]. Less than 10 cases of this entity have been reported [2-5]. We report a case of this unusual tumor in a 4-month-old girl who had lethargy and macrocephaly. Although the clinical, radiographic, and angiographic manifestations of intracranial sarcomas have been described [1-10], the computed tomographic appearance of a primary malignant gliosarcoma has not been published.

Case Report

A 4-month-old girl was noted to have a large head at birth. At age 16 weeks she had lethargy and bulging anterior fontanelle. She was the product of an uneventful 38 week gestation. Her mother (para 1, gravida 1) denied the use of drugs, ethanol, or cigarettes during the pregnancy.

The physical examination failed to reveal any focal neurologic deficits. The head circumference was 46 cm, which far exceeds the 95th percentile for her age. At birth her head measured 35.5 cm, which is at the 90th percentile level.

Unenhanced CT (fig. 1A) demonstrated a large cystic left hemispheric mass obliterating the left lateral ventricle, shifting the midline structures to the right, with moderate dilatation of the right lateral ventricle secondary to obstruction at the foramen of Monroe. The cyst fluid measured 10-12 Hounsfield units (H) above the cerebrospinal fluid values. The cyst was surrounded by focal regions of increased density secondary to calcification in tumor nodules. With contrast administration (fig. 1B), the entire cyst wall showed prominent, nonhomogeneous enhancement. The enhancing margins appeared thin and delicate medially and quite thick laterally and superiorly.

The CT was the only radiographic study required preoperatively. At surgery a large cystic mass replacing most of the left cerebral hemisphere was found. The cyst wall had a white granular appearance. The cyst contents contained viscous, brown, proteinaceous fluid.

Microscopically the tumor was highly cellular and extremely pleomorphic. In some areas it was composed of closely packed,

spindle-shaped cells with small nuclei and bipolar tapering processes. Abundant reticulin fibers were present around tumor cells in these foci (fig. 1C). Other areas contained sheets of variable-size cells with prominent eosinophilic hyaline cytoplasm (fig. 1D). Many




Fig. 1.—A, Noncontrast scan. Large cystic mass in left hemisphere effaces left lateral ventricle and causes shift of midline structures. Focal regions of calcifications within cyst wall (arrows). B, Contrast scan. Thick, nonhomogeneous rim of enhancement surrounds cyst. C, Abundant interconnecting reticulin fiber network surrounds individual tumor cells and several smaller blood vessels. Reticulin stain $\times 370$. D, Sheets of tumor cell with variable size nuclei and prominent hyaline cytoplasm. Many of these cells contained glial fibrillary acidic protein characteristic of gliosarcoma. H and E $\times 600$.

Received October 28, 1980; accepted after revision March 1, 1981.

¹Department of Diagnostic Radiology, University of Kentucky School of Medicine, 800 Rose St., Lexington, KY 40536. Address reprint requests to S. J. Goldstein (room HX315C).

²Department of Surgery, Division of Neurosurgery, University of Kentucky School of Medicine, Lexington, KY 40536.

³Department of Pathology, University of Kentucky School of Medicine, Lexington, KY 40536.

of these cells contained glial fibrillary acidic protein. The neoplasm also showed abundant mitotic figures, small and large foci of necrosis, microcalcification, and a few foci of endothelial proliferation.

Discussion

Malignant gliosarcoma is a truly rare neoplasm previously reported only in adults. It usually spreads by infiltration of the contiguous brain and has a very poor prognosis. Histologically this tumor demonstrates abundant reticular fibers and multiple mitotic figures. Unlike a glioblastoma with sarcomatous elements, the tumor contains little collagen and has a paucity of blood vessels [1, 2].

The CT appearance of this sarcoma in our patient is quite striking. The tumor surrounding the cyst before contrast administration is hyperdense secondary to areas of microcalcification demonstrated histologically. The marked enhancement of the tumor after contrast administration proved to be a reliable indicator of its malignancy [11, 12].

This case is of further interest in that it almost certainly represents a true congenital intracranial neoplasm, as the patient's head was large at birth. In a review of the literature, Solitare and Krigman [13] found only 26 cases of congenital brain tumors that fulfilled their criteria of producing signs or symptoms at birth. Only one congenital cerebral sarcoma has been reported to date and this was an angiosarcoma and not a malignant gliosarcoma as in our case [14].

With the widespread use of CT, most primary intracranial neoplasms in children can be readily detected without the morbidity associated with angiography and pneumoencephalography. Although histopathologic diagnosis is not possible with CT alone, it does provide a safe, noninvasive method for preoperative diagnosis and postoperative follow-up in the neonate and very young child with an intracranial neoplasm.

ACKNOWLEDGMENTS

We thank M. E. Velesio for performing the glial fibrillary acidic protein stains and Glenna S. Williams for secretarial assistance.

REFERENCES

1. Rubinstein LJ. Tumors of the central nervous system. In: *Atlas of tumor pathology*, series 2, fascicle 6. Washington DC: Armed Forces Institute of Pathology, 1972:198-199
2. Rubinstein LJ. The development of contiguous sarcomatous and gliomatous tissue in intracranial tumors. *J Pathol Bacteriol* 1956;71:441-459
3. Kernohan JW, Uihlein A. *Sarcomas of the brain*. Springfield, IL: Thomas, 1962
4. Rubinstein LJ. Morphological problems of brain tumors with mixed cell population. *Acta Neurochir [Suppl] (Wien)* 1964;10:141-158
5. Henry JM, Leestma JE. Astrocytoma arising in meningeal fibrosarcoma. *Acta Neuropathol (Berl)* 1973;23:334-337
6. Fessard C. Cerebral tumors in infancy. *Am J Dis Child* 1968;115:302-308
7. Baker LB. Intracranial sarcomas: their radiographic manifestations. *AJR* 1960;84:70-77
8. Abbot KH, Kernohan JW. Primary sarcomas of the brain: review of the literature and report of twelve cases. *Arch Neurol Psychiatr* 1942;50:43-66
9. Hsu KY. Primary intracranial sarcomas. *Arch Neurol Psychiatr* 1940;43:901-924
10. Liebeskind B, Cohen S, Anderson R, et al. Unusual segmental cerebrovascular changes. *Radiology* 1973;106:119-122
11. Butler AR, Horii SC, Kriceff II, et al. Computed tomography in astrocytomas: a statistical analysis of the parameters of malignancy and the positive contrast-enhanced CT scan. *Radiology* 1978;129:433-439
12. Weisberg LA. Cerebral computed tomography in the diagnosis of supratentorial astrocytomas. *Comput Tomogr* 1980;4:87-105
13. Solitare GB, Krigman MR. Congenital intracranial neoplasm. *Neuropathol Exp. Neurol* 1964;23:280-292
14. Corten MH. Über ein Haemangioma sarcomatodes des Gehirns bei einem Neugeborenen. *Frankfort Z Pathol* 1921;suppl 24: 693