



**Providing Choice & Value**

Generic CT and MRI Contrast Agents



**FRESENIUS  
KABI**

**CONTACT REP**

**AJNR**

**Subtle Malformation of the Cochlear Apex  
and Genetic Abnormalities: Beyond the  
"Thorny" Cochlea**

F. D'Arco, A. Biswas, E. Clement, K. Rajput and A.F.  
Juliano





This information is current as  
of July 30, 2025.

*AJNR Am J Neuroradiol* 2023, 44 (1) 79-81

doi: <https://doi.org/10.3174/ajnr.A7746>

<http://www.ajnr.org/content/44/1/79>

# Subtle Malformation of the Cochlear Apex and Genetic Abnormalities: Beyond the “Thorny” Cochlea

 F. D’Arco,  A. Biswas,  E. Clement,  K. Rajput, and  A.F. Juliano

## ABSTRACT

**SUMMARY:** With the routine use of high-resolution heavily T2-weighted sequences to evaluate patients with hearing deficits, new, subtle phenotypes of cochlear malformations are being discovered and an increasing number of genotype-phenotype correlations are being found through a reverse phenotype approach, which can help guide geneticists. In this brief report, we present subtle malformations of the apical turn of the cochlea related to 3 genetic mutations, emphasizing the importance of a careful assessment of the cochlear apex.

**ABBREVIATIONS:** BOR = branchio-oto-renal syndrome; DEGCAGS = DEvelopmental delay with GIastrointestinal, CArdiovascular, GeNitorinary, and Skeletal abnormalities; IAC = internal auditory canal; SNHL = sensorineural hearing loss

With the use of high-resolution sequences in MR imaging assessment of patients with congenital sensorineural hearing loss (SNHL), newer phenotypes of cochlear malformations have been described, including several types of cochlear hypoplasias.<sup>1,2</sup> There is increasing recognition that specific radiologic appearances of the cochlea and temporal bone may be associated with specific syndromes or genetic mutations.<sup>3-6</sup> Thus, an imaging phenotype can suggest an underlying genotype, contributing to clinical and genetic work-up of the patient.

The normal cochlea has an apical turn that is smooth in contour and evenly tubular throughout its length. The apex (following the basal and middle turns) typically spans 180°–270° around the center point, resulting in a cochlea with 2.5–2.75 turns total.<sup>7</sup>

Here, we describe subtle malformations of the apical aspect of the cochlea in 5 patients with 3 different genetic abnormalities and cochleovestibular symptoms; these apical malformations were chosen among the authors by consensus. The patients were selected from the Great Ormond Street Hospital database of dysplastic cochleae. We retrospectively reviewed the database for all cases that involved an anomaly in the apical

turn, yielding these 5 cases. All these cases had genetic diagnoses available. We describe possible genetic mechanisms that may cause apical turn anomalies, and emphasize the importance of not overlooking such subtle cochlear abnormalities.

## CASE SERIES

### TKFC-Related Disorder

A 7-year-old patient known to have a biallelic mutation in the *TKFC* gene (Mendelian Inheritance in Man, 618805), with congenital cataracts, microphthalmia, and developmental delay, underwent MR imaging of the brain and inner ear/internal auditory canal (IAC) for balance difficulties and hypersensitivity to loud and sudden noises without hearing loss.

MR imaging of the brain demonstrated bilateral cataracts and cerebellar atrophy. A high-resolution 3D heavily T2-weighted sequence of the IAC revealed symmetric subtle cochlear abnormality characterized by a peculiar appearance of the upper part of the cochlea with an extra half-turn beyond the apical turn, resulting in >3 turns total (Fig 1A).<sup>7</sup> The apical turn and the extra half-turn were not pointed but were smooth and flat like a normal apical turn. The cochlear height, measured in accordance with described methods in the literature (Fig 1B), was 6 mm (above the normal range of 4.3–5.4 mm).<sup>8,9</sup> The cochleovestibular nerves were present and normal in course and caliber. The other inner ear structures were normal as well, including preserved internal partitioning of both cochleae.

### Presumed DEGCAGS Syndrome (*ZNF699* Gene)


A 7-month-old boy with a homozygous variant of unknown significance in the *ZNF699* gene (presumed diagnosis of DEvelopmental

Received August 11, 2022; accepted after revision November 15.


From the Departments of Radiology (F.D., A.B.), Clinical Genetics (E.C.), and Audiological Medicine (K.R.), Great Ormond Street Hospital for Children, National Health Service Foundation Trust, London, UK; and Department of Radiology (A.F.J.), Massachusetts Eye and Ear, Harvard Medical School, Boston, Massachusetts.

Please address correspondence to Amy F. Juliano, MD, Department of Radiology, Massachusetts Eye and Ear, 243 Charles St, Boston, MA 02114; e-mail: amy\_juliano@meei.harvard.edu; @amyfjuliano


<http://dx.doi.org/10.3174/ajnr.A7746>



**FIG 1.** A, High-resolution 3D heavily T2-weighted MR images of the inner ear and IAC in a patient with a *TKFC*-related disorder. There is an extra turn (arrows) at the apical portion of the cochlea beyond the usual apical turn and an overall greater cochlear height. The cochlear nerves are present, the modiolus is normal in appearance, and there is normal internal partitioning. Contrast this appearance with that of the normal cochlear apical turn seen in Fig 3B, without an extra turn beyond it. B, Coronal CT image through the cochlea shows the method to measure cochlear height as defined by Shim et al.<sup>8</sup> The maximal height of the cochlea is measured along an axis perpendicular to the oval window (white line).



**FIG 2.** High-resolution 3D heavily T2-weighted MR images of the inner ear and IAC in a patient with a *ZNF699* gene mutation. The uppermost turn of the cochlear apex following the middle turn has a short and stumpy shape (arrows).



**FIG 3.** High-resolution 3D heavily T2-weighted MR image of a typical *SIX1*-BOR protuberant "thorny" tip of the cochlea (arrow in A) in comparison with a healthy control in whom the apical turn is uniform and relatively flat and tubular (arrow in B).

delay with Gastrointestinal, Cardiovascular, Genitourinary, and Skeletal abnormalities [DEGCAGS] syndrome [Mendelian Inheritance in Man, 619488]) presented with bilateral profound SNHL. He also had developmental delay, atrial septal defect, dysmorphic features, and congenital clavicle deformity.

MR imaging of the IAC was performed to investigate the hearing loss. It demonstrated bilateral abnormality of the apical turn of the cochlea, very similar to the "thorny" cochlea described by Pao et al,<sup>10</sup> in relation to *SIX1* mutation in branchio-oto-renal syndrome (BOR). The cochlear apex was irregular, with a short, stumplike shape (Fig 2). There was mild hypoplasia of the cochlear nerve on the left, but otherwise the inner ear structures were normal bilaterally. The cochlear height was 5.8 and 5.7 mm on the right and left, respectively.

### SIX1-Related BOR

Searching our institutional archive for genetic malformations, we found, among patients with a genetic diagnosis of BOR syndrome, 3 patients with confirmed *SIX1* mutation. Five of these 6 cochleae showed a characteristic "thorny" apical turn, with a small, pointed protuberant appearance (Fig 3). These patients presented with SNHL and other clinical abnormalities expected in BOR due to a *SIX1* genetic mutation (ie, otic and branchial defects consistently seen, renal defects variably seen, and less prevalent than among individuals with BOR related to an *EYA1* genetic muta-

tion).<sup>11</sup> Two of them were previously included in the study by Pao et al.<sup>10</sup>

### DISCUSSION

All patients included in this case series presented with labyrinthine symptoms (4 with SNHL and 1 with balance difficulties and sound hypersensitivity) and very subtle malformations of the cochlear apical turn, which were overlooked by the reporting radiologists in 3 of the 5 cases.

The genetics of inner ear development are not completely understood, but we know that, at some point during embryologic development, genes polarize toward the cochlear (ventral) or vestibulocanicular (dorsal) components of the otic capsule.<sup>12</sup> It is also clear, from radiologic studies of cochleae fitting the description of the so-called cochlear hypoplasia type 4 anomaly, that the basal turn of the cochlea may be relatively preserved while the middle and apical turns are more affected.<sup>13</sup> Indeed, this is the case in the "unwound" cochlea in patients with BOR<sup>14,15</sup> and in the extreme hypoplastic cochlea in patients with Walker-Warburg syndrome.<sup>4</sup>

In patients with BOR, both *EYA1* and *SIX1* are expressed in the ventral (cochlear) part of the developing otic vesicle, but the expression of *SIX1* is dependent on that of *EYA1*, and its expression is more prominent in the cochlear apex in murine models.<sup>10,16</sup> This accounts for the different phenotypes seen among patients with BOR with mutations in the *EYA1* versus the *SIX1* gene, with only the apex being malformed in those with *SIX1* mutations.

Although the role of the *TKFC* gene in ear development has not been reported, patients with this mutation may have progressive low-frequency hearing impairment.<sup>17</sup> This specific type of hearing loss may be explained by morphologic abnormalities in the cochlear apex, which allows hearing of low pitches. In fact, the role of the shape of the apical cochlea has been correlated, in a comparative study among different species, with low-frequency hearing limits.<sup>18</sup> Our case of *TKFC*-related disorder represents an example of clinicoradiologic correlation that can shed light on previously unknown functions of specific genes in ear development, including rare genetic mutations that lead to multisystemic abnormalities.

DEGCAGS syndrome is due to an autosomal recessive mutation in the *ZNF699* gene and is characterized by neurodevelopmental

delay, abnormal facial features, growth delay, syndactyly/polydactyly, and anemia/pancytopenia.<sup>19</sup> This gene encodes for a nuclear zinc-finger protein with a possible function in nucleic acid binding.<sup>20</sup> Little is known about the role of this gene, but patients described with this syndrome have SNHL as a constant symptom.<sup>19</sup> It is, therefore, likely that the cause of the SNHL is related to a specific role of this gene in the development of the apical part of the cochlea. Again, the reverse phenotyping approach in this case can help define as yet suboptimally understood gene functions.

## CONCLUSIONS

Subtle abnormalities of the inner ears, particularly those involving the apical part of the cochlea, can be easily overlooked, even with the use of optimal 3D high-resolution heavily-T2-weighted sequences on a 3T MR imaging scanner. We described 5 patients with 3 genetic abnormalities, all characterized by labyrinthine symptoms and malformation of the apical aspect of the cochlea and otherwise normal inner ear anatomy. These findings correlate with clinical symptoms of hearing impairment and shed light on the role of rare genetic abnormalities in inner ear development. A diligent search for these at-times subtle findings must be conducted, especially in cases of MR imaging of the IAC with seemingly negative findings in symptomatic patients.

This report adds to the growing evidence of genotype-phenotype correlation between syndromic and/or congenital deafness and anomaly of the cochlear apex in children.

**Disclosure forms** provided by the authors are available with the full text and PDF of this article at [www.ajnr.org](http://www.ajnr.org).

## REFERENCES

1. Sennaroglu L. **Histopathology of inner ear malformations: do we have enough evidence to explain pathophysiology?** *Cochlear Implants Int* 2016;17:3–20 [CrossRef Medline](#)
2. Quirk B, Youssef A, Ganau M, et al. **Radiological diagnosis of the inner ear malformations in children with sensorineural hearing loss.** *BJR Open* 2019;1:20180050 [CrossRef Medline](#)
3. Siddiqui A, D'Amico A, Colafati GS, et al. **Hypothalamic malformations in patients with X-linked deafness and incomplete partition type 3.** *Neuroradiology* 2019;61:949–52 [CrossRef Medline](#)
4. Talenti G, Robson C, Severino MS, et al. **Characteristic cochlear hypoplasia in patients with Walker-Warburg syndrome: a radiologic study of the inner ear in  $\alpha$ -dystroglycan-related muscular disorders.** *AJNR Am J Neuroradiol* 2021;42:167–72 [CrossRef Medline](#)
5. D'Arco F, Sanverdi E, O'Brien WT, et al. **The link between inner ear malformations and the rest of the body: what we know so far about genetic, imaging and histology.** *Neuroradiology* 2020;62:539–44 [CrossRef Medline](#)
6. D'Arco F, Youssef A, Ioannidou E, et al. **Temporal bone and intracranial abnormalities in syndromic causes of hearing loss: an updated guide.** *Eur J Radiol* 2020;123:108803 [CrossRef Medline](#)
7. Biedron S, Westhofen M, Ilgner J. **On the number of turns in human cochleae.** *Otol Neurotol* 2009;30:414–17 [CrossRef Medline](#)
8. Shim HJ, Shin JE, Chung JW, et al. **Inner ear anomalies in cochlear implantees: importance of radiologic measurements in the classification.** *Otol Neurotol* 2006;27:831–37 [CrossRef Medline](#)
9. D'Arco F, Talenti G, Lakshmanan R, et al. **Do measurements of inner ear structures help in the diagnosis of inner ear malformations? A review of literature.** *Otol Neurotol* 2017;38:e384–92 [CrossRef Medline](#)
10. Pao J, D'Arco F, Clement E, et al. **Re-examining the cochlea in branchio-oto-renal syndrome: genotype-phenotype correlation.** *AJNR Am J Neuroradiol* 2022;43:309–14 [CrossRef Medline](#)
11. Ruf RG, Xu PX, Silvius D, et al. **SIX1 mutations cause branchio-oto-renal syndrome by disruption of EYA1-SIX1-DNA complexes.** *Proc Natl Acad Sci U S A* 2004;101:8090–95 [CrossRef Medline](#)
12. Chatterjee S, Kraus P, Lufkin T. **A symphony of inner ear developmental control genes.** *BMC Genet* 2010;11:68 [CrossRef Medline](#)
13. Talenti G, Manara R, Brotto D, et al. **High-resolution 3 T magnetic resonance findings in cochlear hypoplasias and incomplete partition anomalies: a pictorial essay.** *Br J Radiol* 2018;91:20180120 [CrossRef Medline](#)
14. Hsu A, Desai N, Paldino MJ. **The unwound cochlea: a specific imaging marker of branchio-oto-renal syndrome.** *AJNR Am J Neuroradiol* 2018;39:2345–49 [CrossRef Medline](#)
15. Robson CD. **Congenital hearing impairment.** *Pediatr Radiol* 2006;36:309–24 [CrossRef Medline](#)
16. Zheng W, Huang L, Wei ZB, et al. **The role of Six1 in mammalian auditory system development.** *Development* 2003;130:3989–4000 [CrossRef Medline](#)
17. Wortmann SB, Meunier B, Mestek-Boukhibar L, et al. **Bi-allelic variants in TKFC encoding triokinase/FMN cyclase are associated with cataracts and multisystem disease.** *Am J Hum Genet* 2020;106:256–63 [CrossRef Medline](#)
18. Manoussaki D, Chadwick RS, Ketten DR, et al. **The influence of cochlear shape on low-frequency hearing.** *Proc Natl Acad Sci U S A* 2008;105:6162–66 [CrossRef Medline](#)
19. Bertoli-Avella AM, Kandaswamy KK, Khan S, et al. **Combining exome/genome sequencing with data repository analysis reveals novel gene-disease associations for a wide range of genetic disorders.** *Genet Med* 2021;23:1551–68 [CrossRef Medline](#)
20. Scholz H, Franz M, Heberlein U. **The hangover gene defines a stress pathway required for ethanol tolerance development.** *Nature* 2005;436:845–47 [CrossRef Medline](#)