

# Generic Contrast Agents

Our portfolio is growing to serve you better. Now you have a *choice*.



FRESENIUS  
KABI

[VIEW CATALOG](#)

# AJNR

## Myth of the mesoderm.

A S Smith

*AJNR Am J Neuroradiol* 1989, 10 (2) 449

<http://www.ajnr.org/content/10/2/449.citation>

This information is current as  
of May 9, 2025.

### Myth of the Mesoderm

I would like to clarify a misconception about the origin and content of dermoid vs epidermoid cysts. Reference was made to this in the article "Temporoparietal Dermoid Cysts with Intracranial Extension" by Currarino and Rutledge [1]. Although these authors' findings of dermoid cyst with sinus in an unexpected location are certainly of interest, I must differ with the statement that inclusion of ectodermal and mesodermal elements constitutes a dermoid cyst. This same concept has been repeated in other radiologic reports and major neuroradiologic textbooks [2–5]. The misconception common to these writings is that hair and sebaceous and sweat glands are of mesodermal origin. Although these elements lie within mesodermal connective tissue, the origin of hair follicles and these glandular structures is the ectoderm [6–8]. The embryologic and pathologic literature appears to have no debate about the theories of origin of epidermoid and dermoid cysts from dermis [9–11]. Inclusion of small amounts of mesodermal or endodermal elements in a tumor composed primarily of tissue of ectodermal origin results in its classification as a teratoid tumor rather than a teratoma, in which larger volumes of these additional elements are present [12].

This issue is raised to avoid misinterpretation for the explanation of variations in signal intensity between epidermoid and dermoid cysts by MR. Although increased signal intensity has been seen on T1-weighted images in dermoid and, occasionally, epidermoid cysts, this does not imply the presence of mesodermal fat as contained in lipomas [13]. Desquamated debris of squamous epithelial cells, which contains cholesterol and keratin, is common to both epidermoid and dermoid cysts. In dermoid cysts, the additional breakdown products of hair combined with the secretions of sweat and sebaceous glands result in an oily mixture containing lipid metabolites [12], more fluid than that seen in epidermoid cysts [13]. The calcifications that can be seen in association with a dermoid cyst does not change the classification of this entity as an ectodermal derivation. These calcifications may be dystrophic or a result of dental enamel, another ectodermal derivative [9].

Clarification of the "myth of the mesoderm" of dermoid cyst origin not only is of academic interest but also emphasizes a need to look outside the radiologic literature when making statements about non-radiologic fields, such as embryology and pathology.

Alison S. Smith  
Cleveland Metropolitan General Hospital  
Cleveland, OH 44109

### REFERENCES

1. Currarino G, Rutledge JC. Temporoparietal dermoid cysts with intracranial extension. *AJNR* 1988;9:385–387
2. Ford K, Drayer B, Osborne D, Dubois P. Transient cerebral ischemia as a manifestation of ruptured intracranial dermoid cyst. *J Comput Assist Tomogr* 1981;5(6):895–897
3. Williams AL. Tumors. In: Williams AL, Haughton VM, eds. *Cranial computed tomography: a comprehensive text*. St. Louis: Mosby, 1985:214
4. Lee SH, Rao KCVG. Primary tumors in adults. In: Lee SH, Rao KCVG, eds. *Cranial computed tomography and MRI*. New York: McGraw-Hill, 1983:318–322
5. Roppolo HMN. Intracellar and parasellar abnormalities. In: Latchaw RE, ed. *Computed tomography of the head, neck and spine*. Chicago: Year Book Medical, 1985:343
6. Bloom W, Fawcett DW. *A textbook of histology*, 10th ed. Philadelphia: Saunders, 1975:578.
7. Murphy GF, Kwan TH, Mihm MC. The skin. In: Robin SL, Cotran RS, Kumar V, eds. *Pathologic basis of disease*, 3rd ed. Philadelphia: Saunders, 1984:1257–1303
8. Junqueira LC, Carneiro J, Long JA. *Basic histology*, 5th ed. Los Altos, CA: Lang, 1986:389–412
9. Arey LB. *Developmental anatomy*. Philadelphia: Saunders, 1928:89, 230–235
10. Leech RW, Olafson RA. Epithelial cysts of the neural axis. *Arch Pathol Lab Med* 1977;101:196–202
11. Burger PC, Vogel FS. *Surgical pathology of the nervous system and its coverings*. New York: Wiley & Sons, 1982:117–122
12. Rubinstein LJ. *Atlas of tumor pathology*, fasc. 6: Tumors of the central nervous system. Washington, DC: Armed Forces Institute of Pathology, 1972:285–292
13. Hahn FJ, Ong E, McComb RD, et al. MR imaging of ruptured intracranial dermoid. *J Comput Assist Tomogr* 1986;10(5):888–892

### Reply

We too believe, as does Dr. Smith, that dermoid cysts are ectodermal in origin, but we erroneously chose to follow the concept as established in print. We are grateful that Dr. Smith has clarified the origin of the dermoid cyst for all of us.

Joe C. Rutledge  
Children's Hospital and Medical Center  
Seattle, WA 98105

Guido Currarino  
Children's Medical Center at Southwestern Medical Center  
Dallas, TX 75235

Letters are published at the discretion of the Editor and are subject to editing.  
Letters to the Editor must not be more than two *double-spaced*, typewritten pages. One or two figures may be included. Abbreviations should not be used.  
Material being submitted or published elsewhere should not be duplicated in letters, and authors of letters must disclose financial associations or other possible conflicts of interest.  
Letters concerning a paper published in the *AJNR* will be sent to the authors of the paper for a reply to be published in the same issue. Opinions expressed in the Letters to the Editor do not necessarily reflect the opinions of the Editor.