

Discover Generics

Cost-Effective CT & MRI Contrast Agents



AJNR

Necrotizing myelopathy.

R C Kim

AJNR Am J Neuroradiol 1991, 12 (6) 1084-1086 http://www.ajnr.org/content/12/6/1084.citation

This information is current as of June 5, 2025.

Commentary

Necrotizing Myelopathy

Ronald C. Kim¹

The preceding paper by Mirich et al. [1] describes the MR findings in four subjects with biopsy-proved necrotizing myelopathy. In the first three patients, the clinical course was characterized by progressive motor and sensory dysfunction in the lower extremities, evolving over periods ranging from 3 months to 2 years. In the fourth case, the clinical course was characterized by a Brown-Séquard syndrome with a C4 sensory level that developed over a 3-week period. All four patients had abrupt worsening of their signs and symptoms. In each instance, initial MR imaging showed focal intramedullary expansion of the spinal cord, and histologic examination of the specimen obtained at surgery showed parenchymal coagulative necrosis and thickening and hyalinization of the vascular walls. A definite etiologic diagnosis could not be made.

The development of clinically progressive necrosis of the spinal cord not related to traumatic injury, an infectious or a primary inflammatory process, or infiltration or compression by neoplasm, though not particularly common, may pose difficulties in diagnosis. The purpose of this discussion is to review briefly some of the disorders that might be considered.

Among the circulatory disorders, the most frequently discussed are those that are related to impairment of spinal venous outflow. Clinically, nonhemorrhagic venous infarction of the spinal cord tends to evolve over a period of weeks to months and is typically accompanied by extensive thrombosis within leptomeningeal veins [2]. The longitudinal extent of the damage is variable but may be considerable. Factors that predispose its development include hypercoagulability (e.g., migratory thrombophlebitis or polycythemia rubra vera) or spinal venous hypertension (such as that seen in the presence

of a spinal dural arteriovenous fistula [AVF]. The existence of AVFs draining into the spinal leptomeningeal venous system was recognized only relatively recently, with the aid of selective spinal angiography [3] and angiotomography [4, 5], but these AVFs are now thought to be the commonest type of spinal vascular malformation [6-9]. Although the feeding arteries may be situated outside the spinal canal [4, 10], in most instances the vascular nidus is embedded within or is adjacent to the dura ensheathing the proximal spinal nerve root [7]. thus giving rise to the term spinal dural AVF. Most of these lesions are found at or below midthoracic levels [4, 6-8]. Despite the absence in the literature of any descriptions of the pathologic findings within the spinal cord of angiographically or pathologically confirmed spinal dural AVFs, it is widely thought that the alterations correspond to those described initially by Foix and Alajouanine [11] and subsequently by others [12-15]. These were given a variety of names, including subacute necrotic myelitis, Foix-Alajouanine syndrome, angioma racemosum venosum, angiodysgenetic necrotizing myelopathy, type I spinal arteriovenous malformation (AVM), thrombosing spinal AVM, and spinal AVM with acute neurologic deterioration [11, 13, 16-19]. The general belief is that shunting of arterial blood into spinal leptomeningeal veins results in venous hypertension, which in turn leads to interference with venous outflow from the spinal cord, progressive intramedullary congestion, mural thickening and hyalinization within the venocapillary network, and hypoxic parenchymal atrophy [4, 7, 15].

Necrotizing myelopathy is also a feature of the condition referred to as neuromyelitis optica or Devic's syndrome. Although the symptom complex classically consists of a com-

AJNR 12:1084-1086, November/December 1991; AJR 158: January 1992 0195-6108/91/1206-1084 © American Society of Neuroradiology

This article is a commentary on the preceding article by Mirich et al.

¹ Laboratory Service (113NP), Veterans Affairs Medical Center, Long Beach, CA 90822, and the Department of Pathology, University of California, Irvine, Irvine, CA 92717. Address reprint requests to R. C. Kim at the Veterans Affairs Medical Center.

bination of blindness and paraplegia, there is no reason why the many disorders that have been associated with it, including multiple sclerosis [20, 21], acute disseminated encephalomyelitis [21], varicella-zoster infection [22, 23], mumps [23], rubeola [23], infectious mononucleosis [24], systemic lupus erythematosus (SLE) [25, 26], pulmonary tuberculosis [27-29], and clioquinol intoxication [30-32], could not lead to the development of myelopathy alone. It has been repeatedly emphasized, for example, that the spinal cord lesions in Japanese patients with multiple sclerosis tend to be more necrotic than the lesions in other patients [33, 34]. Additionally, of the three cases of noninflammatory necrotizing myelopathy described by Hughes and Mair [27] in patients with pulmonary tuberculosis, only one had evidence of involvement of the optic nerve. Finally, lupus myelopathy in the absence of optic neuropathy has been recognized for many years [35-38].

The subject of lupus myelopathy deserves further comment. Although myelopathy usually appears within a few years of the time at which a diagnosis of SLE is made, in a minority (15-20%) it is the presenting feature [38]. Typically, the course of illness is one of fairly rapid progression, but at times it is characterized by remissions and relapses. In most instances, the spinal cord is affected at or below thoracic levels, but approximately one quarter of the cases have involved the cervical region. Pathologically the lesion consists of one or more foci of subtotal coagulative or liquefactive necrosis [35, 37, 39, 40], although, occasionally, circumferentially distributed pallor of the white matter with ballooning degeneration of myelin sheaths and disruption of axons has been described [36, 40]. The pathogenesis of the myelopathy has not been determined, but at least two groups of investigators have recently suggested a relationship to the presence of antiphospholipid antibodies [41, 42].

The existence of a malignancy-associated or paraneoplastic necrotizing myelopathy not associated with radiation therapy or with compression or infiltration by tumor was first suggested by Mancall and Rosales in 1964 [43]. The subject has been reviewed periodically since then [44-48]. The typical clinical course is one of fairly rapidly progressive neurologic dysfunction that reflects involvement of the spinal cord at any level. The condition, which has been described in association with a variety of neoplasms (most notably bronchopulmonary and lymphoreticular malignancies), is characterized by massive or patchy multifocal coagulative necrosis affecting both gray and white matter, with a paucity of inflammatory cells. The pathogenesis of the disorder is unknown. Recently, however, Iwamasa et al. [49] have described two cases of malignancy-associated necrotizing myelopathy in which, despite the relative scarcity of inflammatory cells, the presence of herpes simplex virus type 2 was demonstrated both immunohistochemically and by electron microscopy. This raises the possibility that the cause of some of the other noninflammatory necrotizing myelopathies that have been included in this discussion may, at times, be viral.

The last major form of noninflammatory necrotizing myelopathy is not associated with any other pathologic disorder. This condition, which was called "acute necrotic myelopathy" by Hoffman [50] and "progressive necrotic myelopathy" by Folliss and Netsky [51], is entirely idiopathic. The pathologic findings, which may vary in their extent, are those of coagulative or liquefactive necrosis in the absence of striking infiltration by inflammatory cells. It is likely that, as knowledge advances, this form of myelopathy will be shown to be etiologically heterogeneous.

The confusion caused by the noninflammatory necrotizing myelopathies is attributable to our lack of knowledge of the pathogenesis of most of these disorders, the difficulty of access to adequate amounts of tissue for diagnostic purposes, the relatively small number of persons with expertise in spinal angiography, the lack of suitable radiologic methods for the study of the spinal parenchymal and spinal leptomeningeal venous systems, and the relatively low frequency with which the spinal cord is examined at autopsy.

In the patients described by Mirich et al. [1], we can only speculate about the cause of the necrotizing myelopathy. Although selective spinal angiography was not done, it is certainly possible that some of them may have had spinal dural AVFs. The finding of thickening and hyalinization of vascular walls is not specific, however, and may have been caused by the spinal cord necrosis [51]. More suggestive would have been the presence of marked thickening of the walls of leptomeningeal veins. Moreover, involvement of the cervical spinal cord, as was seen in case 4, is most unusual, and the development of a Brown-Séquard syndrome has not, to my knowledge, previously been described in association with a spinal dural AVF. It is likely, therefore, that more than one disorder is represented in this interesting group of patients, and I hope that MR imaging studies of the type described by Mirich et al. will be beneficial to our understanding of this mystifying form of spinal cord disease.

REFERENCES

- Mirich DR, Kucharczyk W, Keller MA, Deck J. Subacute necrotizing myelopathy: MR imaging in four pathologically proved cases. *AJNR* 1991;12:1077–1083
- Kim RC, Smith HR, Henbest ML, Choi BH. Nonhemorrhagic venous infarction of the spinal cord. Ann Neurol 1984;15:379–385
- Kendall BE, Logue V. Spinal epidural angiomatous malformations draining into intrathecal veins. *Neuroradiology* 1977;13:181–189
- Merland JJ, Riche MC, Chiras J. Les fistules artério-veineuses intracanalaires, extra-médullaires à drainage veineux médullaire. J. Neuroradiol 1980;7:271–320
- Gueguen B, Merland JJ, Riche MC, Rey A. Vascular malformations of the spinal cord: intrathecal perimedullary arteriovenous fistulas fed by medullary arteries. *Neurology* **1987**;37:969–979
- Symon L, Kuyama H, Kendall B. Dural arteriovenous malformations of the spine: clinical features and surgical results in 55 cases. *J Neurosurg* 1984;60:238–247
- Rosenblum B, Oldfield EH, Doppman JL, DiChiro G. Spinal arteriovenous malformations: a comparison of dural arteriovenous fistulas and intradural AVM's in 81 patients. *J Neurosurg* 1987;67:795–802
- Thron AK. Vascular anatomy of the spinal cord: neuroradiological investigations and clinical syndromes. New York: Springer-Verlag, 1988
- Koenig E, Thron A, Schrader V, Dichgans J. Spinal arteriovenous malformations and fistulae: clinical, neuroradiological and neurophysiological findings. J Neurol 1989;236:260–266
- Wrobel CJ, Oldfield EH, DiChiro G, Tarlov EC, Baker RA, Doppman JL. Myelopathy due to intracranial dural arteriovenous fistulas draining intrathecally into spinal medullary veins: report of three cases. *J Neurosurg* 1988;69:934–939

- Foix C, Alajouanine T. La myélite nécrotique subaiguë. Rev Neurol 1926;33:1–42
- Greenfield JG, Turner JWA. Acute and subacute necrotic myelitis. Brain 1939;62:227–252
- Wyburn-Mason R. The vascular abnormalities and tumours of the spinal cord and its membranes. London:Kimpton, 1943
- Mair WGP, Folkerts JF. Necrosis of the spinal cord due to thrombophlebitis (subacute necrotic myelitis). *Brain* 1953;76:563–575
- Aminoff MJ, Barnard RO, Logue V. The pathophysiology of spinal vascular malformations. J Neurol Sci 1974;23:255–263
- Scholz W, Manuelidis EE. Myélite nécrotique (Foix-Alajouanine): angiodysgenetische nekrotisierende Myelopathie. *Dtsch Z Nervenheilk* 1851;165:56–71
- 17. Doppman JL, DiChiro G, Ommaya AK. Selective arteriography of the spinal cord. St. Louis: Warren H Green, **1969**
- Wirth FP Jr, Post KD, DiChiro G, Doppman JL, Ommaya AK. Foix-Alajouanine disease: spontaneous thrombosis of a spinal cord arteriovenous malformation: a case report. *Neurology* **1970**;20:1114–1118
- Criscuolo GR, Oldfield EH, Doppman JL. Reversible acute and subacute myelopathy in patients with dural arteriovenous fistulas: Foix-Alajouanine syndrome reconsidered. *J Neurosurg* **1989**;70:354–359
- Stansbury FC. Neuromyelitis optica (Devic's disease): presentation of five cases, with pathologic study, and review of literature. Arch Ophthalmol 1949;42:292–335, 465–501
- Cloys DE, Netsky MG. Neuromyelitis optica. In: Vinken PJ, Bruyn GW, eds. Handbook of clinical neurology, vol. 9. Multiple sclerosis and other demyelinating diseases. Amsterdam: North-Holland, **1970**:426–436
- McAlpine D, Kuroiwa Y, Toyokura Y, Araki S. Acute demyelinating disease complicating herpes zoster. J Neurol Neurosurg Psychiatry 1959;22: 120–123
- Chusid MJ, Williamson SJ, Murphy JV, Ramey LS. Neuromyelitis optica (Devic disease) following varicella infection. J Pediatr 1979;95:737–738
- Williamson PM. Neuromyelitis optica following infectious mononucleosis. Proc Aust Assoc Neurol 1975;12:153–155
- April RS, Vansonnenberg E. A case of neuromyelitis optica (Devic's syndrome) in systemic lupus erythematosus: clinicopathologic report and review of the literature. *Neurology* **1976**;26:1066–1070
- Kinney EL, Berdoff RL, Rao NS, Fox LM. Devic's syndrome and lupus erythematosus: a case report with necropsy. Arch Neurol 1979;36: 643–644
- Hughes RAC, Mair WGP. Acute necrotic myelopathy with pulmonary tuberculosis. *Brain* 1977;100:223–238
- Barbizet J, Degos J-D, Meyrignac CC. Neuromyélite optique aiguë associée à une tuberculose pulmonaire aiguë. *Rev Neurol* 1980;136:303–309
- Silber MH, Willcox PA, Bowen RM, Unger A. Neuromyelitis optica (Devic's syndrome) and pulmonary tuberculosis. *Neurology* **1990**;40:934–938
- Tateishi J, Kuroda S, Saito A, Otsuki S. Experimental myelooptico neuropathy induced by clioquinol. Acta Neuropathol 1973;24:304–320
- Rose FC, Gawel M. Clioquinol neurotoxicity: an overview. Acta Neurol Scand 1984;70[suppl 100]:137–145
- Pallis C. Some thoughts on 'SMON'. Acta Neurol Scand 1984;70[suppl 100]:147–153
- 33. Itoyama Y, Ohnishi A, Tateishi J, Kuroiwa Y, Webster HdeF. Spinal cord

multiple sclerosis lesions in Japanese patients: schwann cell remyelination occurs in areas that lack glial fibrillary acidic protein (GFAP). Acta Neuropathol **1985**;65:217–223

- Yamamoto T, Kawamura J, Hashimoto S, Nakamura M. Extensive proliferation of peripheral type myelin in necrotic spinal cord lesions of multiple sclerosis. *J Neurol Sci* **1991**;102:163–169
- Penn AS, Rowan AJ. Myelopathy in systemic lupus erythematosus. Arch Neurol 1968;18:337–349
- Johnson RT, Richardson EP. The neurological manifestations of systemic lupus erythematosus: a clinical-pathological study of 24 cases and review of the literature. *Medicine* **1968**;47:337–369
- Andrianakos AA, Duffy J, Suzuki M, Sharp JT. Transverse myelopathy in systemic lupus erythematosus: report of three cases and review of the literature. *Ann Intern Med* 1975;83:616–624
- Al-Husaini A, Jamal GA. Myelopathy as the main presenting feature of systemic lupus erythematosus. *Eur Neurol* 1985;24:94–106
- Allen IV, Millar JHD, Kirk J, Shillington RKA. Systemic lupus erythematosus clinically resembling multiple sclerosis and with unusual pathological and ultrastructural features. J Neurol Neurosurg Psychiatry 1979;42:392–401
- Nakano I, Mannen T, Mizutani T, Yokohari R. Peripheral white matter lesions of the spinal cord with changes in small arachnoid arteries in systemic lupus erythematosus. *Clin Neuropathol* **1989**;8:102–108
- Liu GT, Greene JM, Charness ME. Brown-Séquard syndrome in a patient with systemic lupus erythematosus. *Neurology* 1990;40:1474–1475
- Lavalle C, Pizarro S, Drenkard C, Sánchez-Guerrero J, Alarcón-Segovia D. Transverse myelitis: a manifestation of systemic lupus erythematosus strongly associated with antiphospholipid antibodies. *J Rheumatol* 1990;17:34–37
- Mancall EL, Rosales RK. Necrotizing myelopathy associated with visceral carcinoma. *Brain* 1964;87:639–656
- Norris FH Jr. Remote effects of cancer on the spinal cord. In: Vinken PJ, Bruyn GW, Klawans HL, eds. *Handbook of clinical neurology*, vol. 38. *Neurological manifestations of systemic diseases, part I.* Amsterdam: North-Holland, **1979**:669–677
- Sieben GM, De Reuck JL, De Bruyne JC, Vander Eecken HM. Subacute necrotic myelopathy: its appearance eight years after cure of a breast carcinoma. Arch Neurol 1981;38:775–777
- Henson RA, Urich H. Cancer and the nervous system: the neurological manifestations of systemic malignant disease. Oxford, England: Blackwell, 1982
- Handforth A, Nag S, Sharp D, Robertson DM. Paraneoplastic subacute necrotic myelopathy. *Can J Neurol Sci* **1983**;10:204–207
- Ojeda VJ. Necrotizing myelopathy associated with malignancy: a clinicopathologic study of two cases and literature review. *Cancer* 1984;53:1115–1123
- Iwamasa T, Utsumi Y, Sakuda H, et al. Two cases of necrotizing myelopathy associated with malignancy caused by herpes simplex virus type 2. *Acta Neuropathol* **1989**;78:252–257
- 50. Hoffman HL. Acute necrotic myelopathy. Brain 1955;78:377-393
- Folliss AGH, Netsky MG. Progressive necrotic myelopathy. In: Vinken PJ, Bruyn GW, eds. Handbook of clinical neurology, vol. 9. Multiple sclerosis and other demyelinating diseases. Amsterdam: North-Holland, 1970: 452–468