



Discover Generics

Cost-Effective CT & MRI Contrast Agents



FRESENIUS
KABI

WATCH VIDEO

AJNR

Crossed cerebellar diaschisis.

M Reivich

AJNR Am J Neuroradiol 1992, 13 (1) 62-64

<http://www.ajnr.org/content/13/1/62.citation>

This information is current as
of June 21, 2025.

Crossed Cerebellar Diaschisis

Martin Reivich¹

The term "diaschisis," as originally used by Von-Monakow (1), implies an immediate decrease in neuronal activity in a region due to an interruption of its afferent axonal supply. Experimental studies, in animals, of unilateral ischemia in which neuronal function in the contralateral hemisphere was demonstrated to be impaired, provided evidence for such a mechanism (2). The corpus callosum appeared to play a role in this process. Changes in cerebral blood flow in the hemisphere opposite a cerebral infarction were first demonstrated by Kempinsky et al in 1961 (3), and Hoedt-Rasmussen and Skinhoj in 1964 (4). Several theories have been proposed to explain these remote alterations in blood flow, postulating neurogenic, vasogenic, and chemical mechanisms (3-10).

The remote blood flow changes observed, according to the neurogenic theory, are presumed to be secondary to metabolic alterations produced by the decreased axonal input to the region. Such decreased metabolism in the non-ischemic hemisphere and in the contralateral cerebellum of patients with infarcts has been documented (8, 11-13).

Little data is available concerning the time course of these cerebral blood flow changes in the nonischemic hemisphere. A progressive decline in cerebral blood flow in the nonischemic hemisphere has been observed during the first week after an acute infarction (14). This decline in flow could be partially explained by a loss of autoregulation, but suggested a process more complex than destruction of axonal afferents to the nonischemic region.

Vasoactive substances released from ischemic tissue have been postulated to play a role in these remote effects (15). Experimental data have demonstrated the release of vasoactive substances and neurotransmitters from infarcted brain (16).

While the release of such vasoactive substances might play a role in a more generalized depression of flow and metabolism, the more focal effects seen in the cerebellar hemisphere contralateral to various types of lesions make this less likely to be an important factor in the production of crossed cerebellar diaschisis.

Crossed cerebellar diaschisis has been observed not only following cerebral ischemic events (12, 13), but in patients with brain tumors (13, 17) and following the injection of intracarotid sodium Amytal (18).

One of the earliest studies demonstrating the importance of the pathways between the cerebrum and cerebellum on the function of the latter was that of von Monakow in 1885 in which he noted hypoplasia of the contralateral cerebellar hemisphere following experimental cortical ablation (19). In the clinical literature, one of the earliest reports of similar findings was that of Hassin in 1935, in which atrophy of the contralateral cerebellum was observed in a patient with cerebral atrophy secondary to presumed birth injury (20). Ataxic hemiparesis (21) and other ataxic syndromes have been noted secondary to cortical insults (22, 23). False localized limb ataxia has been found to occur contralateral to cerebral tumors located either anteriorly or posteriorly in the cerebral hemispheres (24). Although such syndromes have been considered to occur mainly from contralateral frontal lesions (25), they also occur with lesions limited to the parietal lobe (22).

The cerebellum has a significant input from the cerebral hemispheres, the bulk of which is contained in the cortico-ponto-cerebellar pathway (26). This pathway arises in the motor, premotor, parietal association, and occipital cortices (27). Other cerebro-cerebellar connections are relayed through the inferior olive and reticular formation (28). The contribution of different regions of the cortex to the cortico-ponto-cerebellar pathway varies in different species. Parietal areas project most heavily to the cerebellum in the cat

¹ Director, Cerebrovascular Research Center, University of Pennsylvania, Philadelphia, PA 19104-6063.

Index terms: Cerebral blood flow; Arteries, carotid; Interventional neuroradiology, complications of; Positron emission tomography (PET); Single photon emission computed tomography (SPECT)

(29), whereas in the monkey, the majority of the descending input to the cerebellum is from the motor and premotor areas (30). Connections from the cerebellum to the cerebral cortex are also widespread and include the motor, premotor, and parietal association cortices (31). These ascending pathways are relayed by the thalamus, reticular formation, and the red nucleus (32). Interruption of this anatomic substrate connecting the cerebral hemispheres and the cerebellum provides a likely mechanism for the production of crossed cerebellar hypometabolism, which has been described with such lesions in humans (12, 13, 17). Animal studies have also demonstrated reduced blood flow in the cerebellum of gerbils within hours of acute cerebral ischemia (33). In addition, increased metabolism has been demonstrated in the cerebellum of rats, contralateral to a focus of cortical seizure activity (34).

In humans, the presence of contralateral cerebellar hypometabolism does not appear to be dependent on a single factor, such as the type, size, or severity of the lesion. It is equally likely to occur in cases of tumor and infarction (12). In general, however, widespread cerebral lesions, with marked hypometabolism, are more likely to be accompanied by contralateral cerebellar hypometabolism. Exceptions to this, however, are not uncommon. Relatively small lesions with mild metabolic depression also have been noted to produce contralateral cerebellar hypometabolism. The type of neurologic deficit also does not appear to be related to the presence of contralateral cerebellar diaschisis. Pure sensory syndromes are accompanied by cerebellar hypometabolism and, conversely, this metabolic abnormality may not be present in some patients with hemiplegia. The presence of a parietal lesion, however, does appear to be associated more often with contralateral cerebellar hypometabolism. This may be the pathophysiologic correlate of the ataxia that can arise from lesions of the parietal lobe. It also suggests the prominence of the parietal cortex in the cortico-ponto-cerebellar pathway in humans.

There is little information available concerning the time course of crossed cerebellar diaschisis. In one report of reduced cerebral blood flow and oxygen consumption in the contralateral cerebellum in patients with acute stroke, the effect was not seen in patients studied more than 2 months beyond the ictus (35). Similar findings were observed in another report, in which cerebral blood flow and oxygen metabolism were studied in patients with cerebral infarctions (13). It was also

noted that the most intense cerebellar hypometabolism was produced with parietal infarction. Martin and Raichle (36) studied a patient both less than and more than 3 months after an acute stroke and found that the contralateral cerebellar hypometabolism initially present had disappeared. Two patients restudied more than 3 months after an acute stroke by Kushner et al (12) no longer demonstrated contralateral cerebellar hypometabolism that had been present earlier. The mechanism by which cerebellar hypometabolism returns to normal is uncertain. Restoration of function in reversibly affected cerebral areas is one possibility. A more speculative hypothesis may be an increase in activity in alternate or subordinate cerebro-cerebellar pathways to compensate for the primarily affected cortical area.

The present issue of this journal contains two case reports of crossed cerebellar diaschisis that shed additional light on the time course of this phenomenon (37, 38). Both reports describe the prompt occurrence (within 1 min in one case and 10 sec in the other) of decreased blood flow in the contralateral cerebellum following temporary balloon occlusion of the internal carotid artery in patients being evaluated for the treatment of cerebral aneurysms. These studies confirm the rapidity of development of crossed cerebellar diaschisis previously observed in patients undergoing intracarotid sodium Amytal injection (17, 39, 40). In one report (37), a repeat flow study 15 minutes after deflation of the balloon revealed flow had returned to normal. The prompt onset and resolution of crossed cerebellar diaschisis lends support for the theory that this phenomenon is due to transneuronal loss of excitatory input to the cerebellum and that this functional alteration need not result in pathologic alterations in the cerebellum for crossed cerebellar diaschisis to be manifest. Crossed cerebellar diaschisis can produce clinical findings of cerebellar dysfunction, although this is often difficult to assess, since there is often limb dysfunction associated with the primary lesion. These studies lend support to the concept that improvement in neurologic function associated with cerebral insult may in part be due to the resolution of diaschisis as suggested by Feeney and Baron (41).

References

1. Von-Monakow C. *Die Lokalisation im Grosshirn und der Abbau der Funktion Durch Korrikale Heroe*. Wiesbaden, Germany: JF Bergman, 1914:26-34

2. Kempinsky WH. Experimental study of distant effects of acute focal brain injury. *Arch Neurol Psychiatry* 1958;79:376-389
3. Kempinsky W, Boniface WR, Keating JBA, Morgan PP. Serial hemodynamic study of cerebral infarction in man. *Circ Res* 1961;9:1051-1058
4. Hoedt-Rasmussen K, Skinhoj E. Transneuronal depression of the cerebral hemispheric metabolism in man. *Acta Neurol Scand* 1964;40:41-46
5. Skinhoj E. Bilateral depression of CBF in unilateral cerebral disease. *Acta Neurol Scand* 1965;41(suppl 14):161-163
6. McHenry L. Cerebral blood flow studies in middle cerebral and internal carotid artery occlusion. *Neurology* 1966;16:1145-1151
7. Fieschi C, Agnoli A, Battistini N, Bozzao L. Regional cerebral blood flow in patients with brain infarcts. *Arch Neurol* 1966;15:653-660
8. Meyer JS, Shinohara Y, Kanda T, Fukuuchi Y, Ericsson AD, Kok NK. Diaschisis resulting from acute unilateral cerebral infarction. *Neurology* 1970;23:241-247
9. Fujishima M, Tanaka K, Takeya Y, Omae T. Bilateral reduction of hemispheric blood flow in patients with unilateral cerebral infarction. *Stroke* 1974;5:648-653
10. Lavy S, Eldad M, Portnoy Z. Effect of cerebral infarction on the regional cerebral blood flow of the contralateral hemisphere. *Stroke* 1975;6:160-163
11. Meyer JS, Kanda T, Fukuuchi Y, Shimazu K, Dennis EW, Ericsson AD. Clinical prognosis correlated with hemispheric blood flow in cerebral infarction. *Stroke* 1971;2:383-394
12. Kushner JM, Alavi A, Reivich M, Dann R, Burke A, Robinson G. Contralateral cerebellar hypometabolism following cerebral insult. *Ann Neurol* 1984;15:425-434
13. Lenzi GL, Frackowiak RSJ, Jones T. Cerebral oxygen metabolism and blood flow in human cerebral ischemic infarction. *J Cereb Blood Flow Metab* 1982;2:321-335
14. Slater R, Reivich M, Goldberg H, Banka R, Greenberg J. Diaschisis with cerebral infarction. *Stroke* 1977;8:684-690
15. Meyer JS, Welch KMA, Okamoto S, Shimazu K. Disordered neurotransmitter function. *Brain* 1974;97:655-664
16. Globus MY-T, Busto R, Martinez E, Valdes I, Dietrich WD, Ginsberg MD. Comparative effect of transient global ischemia on extracellular levels of glutamate, glycine, and γ -aminobutyric acid in vulnerable and nonvulnerable brain regions in the rat. *J Neurochem* 1991;57:470-478
17. Patronas NJ, Di Ciro G, Smith BH, et al. Depressed cerebellar glucose metabolism in supratentorial tumors. *Brain Res* 1984;291:93-101
18. Kurthen M, Reichmann K, Linke DB, et al. Crossed cerebellar diaschisis in intracarotid sodium amytal procedures: a SPECT study. *Acta Neurol Scand* 1990;81:416-422
19. Von-Monakow C. Experimentelle und pathologisch-anatomische Untersuchungen über die Haubenregion, den Sehlhugel und die regio subthalamica. *Arch Psychiatr Nervenkr* 1885;27:1
20. Hassin GB. Crossed atrophy of the cerebellum: pathologic study of a case. *Arch Neurol Psychiatry* 1935;33:917-928
21. Fisher CM. Ataxic hemiparesis: a pathologic study. *Arch Neurol* 1978;35:126-238
22. Critchley M. *The parietal lobes*. New York: Hafner, 1953:160-163
23. Gassel MM. False localizing signs. *Arch Neurol* 1964;4:70-98
24. Grant FC. Cerebellar symptoms produced by supratentorial tumor. *Arch Neurol Psychiatry* 1928;20:292-317
25. Brun R. Zur Frage der Stirnhirntaxie ein neues Stirnhirnsymptom. *Z Neurol Psychiatr* 1932;138:122
26. Allen GI, Tuskahara N. Cerebrocerebellar communication systems. *Physiol Rev* 1974;54:957-1006
27. Brodal A. Cerebrocerebellar pathways: anatomical data and some functional implications. *Acta Neurol Scand* 1972;51(suppl):153-195
28. Brodal P. The corticopontine projection in the rhesus monkey: origin and principles of organization. *Brain* 1978;101:251-283
29. Sasaki K, Oka H, Matsuda Y. Electrophysiological studies of the projections from the parietal association area to the cerebellar cortex. *Exp Brain Res* 1975;23:91-102
30. Sasaki K, Oka H, Kawaguchi J. Mossy fibre and climbing fibre responses produced in the cerebellar cortex in monkeys. *Exp Brain Res* 1977;29:419-428
31. Sasaki K, Kawaguchi J, Oka H. Electrophysiological studies on the cerebello-cerebral projections in monkeys. *Exp Brain Res* 1976;24:495-507
32. Evarts EV, Thach WT. Motor mechanisms of the CNS: cerebrocerebellar interrelations. *Ann Rev Physiol* 1969;31:451-498
33. Naritomi H. Transtentorial diaschisis: reduction of cerebellar blood flow caused by supratentorial local cerebral ischemia in the gerbil. *Stroke* 1983;14:213-214
34. Collins RC, Kennedy C, Sokoloff L, Plum F. Metabolic anatomy of focal motor seizures. *Arch Neurol* 1976;33:536-542
35. Baron JC, Bousser MG, Comar D, et al. Crossed cerebellar diaschisis: a remote functional depression secondary to supratentorial infarction in man. *J Cereb Blood Flow Metab* 1981;1(suppl 1):S500-S501
36. Martin WRW, Raichle ME. Cerebellar blood flow and metabolism in cerebral hemispheric infarction. *Ann Neurol* 1983;14:168-176
37. Brunberg JA, Frey KA, Horton JA, Kuhl DE. Crossed cerebellar diaschisis: occurrence and resolution demonstrated with PET during carotid temporary balloon occlusion. *AJNR* 1992;13:58-61
38. Eckard DA, Purdy PD, Bonte F. Crossed cerebellar diaschisis and loss of consciousness during temporary balloon occlusion of the internal carotid artery. *AJNR* 1992;13:55-57
39. Ryding E, Sjöholm H, Skeidsvoll H, et al. Delayed decrease in hemispheric cerebral blood flow during Wada test demonstrated by ^{99m}Tc -HMPAO single photon emission computed tomography. *Acta Neurol Scand* 1989;80:248-254
40. Biersack HJ, Linke D, Brassel F, et al. Technetium- 99m HMPAO brain SPECT in epileptic patients before and during unilateral hemispheric anesthesia (Wada test): report of three cases. *J Nucl Med* 1987;28:1763-1767
41. Feeney DM, Baron JC. Diaschisis. *Stroke* 1986;86:817-830