



Get Clarity On Generics

Cost-Effective CT & MRI Contrast Agents



FRESENIUS
KABI

WATCH VIDEO

AJNR

Intraventricular hemorrhage caused by intraventricular meningioma: CT appearance.

I Lang, A Jackson and F A Strang

AJNR Am J Neuroradiol 1995, 16 (6) 1378-1381

<http://www.ajnr.org/content/16/6/1378.citation>

This information is current as
of August 23, 2025.

Intraventricular Hemorrhage Caused by Intraventricular Meningioma: CT Appearance

I. Lang, A. Jackson, and F. A. Strang

Summary: In a patient with intraventricular meningioma presenting with intraventricular hemorrhage, CT demonstrated a well-defined tumor mass in the trigone of the left lateral ventricle with extensive surrounding hematoma.

Index terms: Meninges, neoplasms; Brain, ventricles; Cerebral hemorrhage

Intraventricular meningioma is a rare tumor with approximately 170 previously reported cases (1–7). Despite this it is the most common intraventricular tumor in adults, in whom it is most often located in the trigone of the lateral ventricle (1, 2). Computed tomography (CT) features are similar to meningiomas elsewhere, and in most cases they appear as well-defined high-attenuation masses that show marked enhancement (1, 2). Presentation with subarachnoid hemorrhage is rare (9). We describe a case of intraventricular hemorrhage attributable to intraventricular meningioma and present the CT appearance.

Case Report

A 64-year-old man was admitted with sudden onset of severe occipital headache associated with loss of consciousness. There was a history of treated hypertension and angina. The patient also gave a history of a severe headache of sudden onset approximately 6 weeks previously. Physical examination demonstrated a dense right hemiparesis, a right homonymous hemianopia, and marked expressive nominative dysphasia. Fundoscopy revealed bilateral papilloedema.

A CT scan (Fig 1) showed a well-defined soft-tissue-attenuation lesion lying within the occipital horn of the left lateral ventricle. This mass was approximately $4 \times 5 \times 4$ cm in size and displaced the choroid plexus posteriorly. The mass showed a rim of high attenuation (120 HU),

which extended into the body of the lateral ventricle, through the foramen of Monro and into the third and fourth ventricles. On postcontrast CT (iohexol 340, 50 mL) the lesion showed homogeneous enhancement (Fig 2).

At surgery, the occipital pole of the left lateral ventricle was opened via a temporoparietal craniotomy. A “hard” avascular tumor surrounded by fresh clot was exposed. The tumor arose from a pedicle attached to the choroid plexus. It was removed without complication, and the associated hematoma was evacuated. Histologic examination confirmed the diagnosis of a fibroblastic meningioma. Postoperatively, there was a good recovery of the left hemiparesis, although there was some residual hemianopia and dysphasia.

Discussion

Meningiomas arising within the ventricular system are rare but well described, constituting approximately 0.5% to 2% of all intracranial meningiomas (4, 6) (Kaplan ES, *Parasagittal Meningiomas: A Clinico-pathological Study*, Rochester, Minn: University of Minnesota; 1964, thesis). In the Mayo Clinic series, Kaplan found only 7 intraventricular meningiomas in a series of 1454. Our survey reveals approximately 170 intraventricular meningiomas previously described in the literature (1–7). They most commonly arise in the trigone of the lateral ventricle as in the current case (1, 2, 5, 8). They are slightly more common on the left and show a female predominance of approximately 2:1. Although there is a wide age range, intraventricular meningiomas are most common over the age of 30 years, and the majority present in the third to the sixth decades (1, 4, 6). Meningiomas rarely may arise in the third ventricle, although this is more common in childhood (10, 11).

Received January 27, 1994; accepted after revision June 2.

From the Departments of Neuroradiology (I.L.) and Neurosurgery (F.A.S.), Manchester Royal Infirmary, Manchester, United Kingdom, and the Department of Diagnostic Radiology (A.J.), Manchester University Medical School, Manchester, United Kingdom.

Address reprint requests to A. Jackson, Department of Diagnostic Radiology, Manchester University Medical School, Manchester, United Kingdom.

AJNR 16:1378–1381, Jun 1995 0195-6108/95/1606–1378 © American Society of Neuroradiology




Fig 1. A, Axial CT scan shows a mass lesion in the trigone of the left lateral ventricle (*long arrow*) with surrounding hemorrhage extending into the lateral ventricle (*short arrow*).

B, CT scan shows a rim of hemorrhage around the tumor (*long arrow*) and hemorrhage extending through the foramen of Monro (*short arrow*).

C, CT scan shows a rim of hemorrhage around the tumor (*long arrow*) and blood in the third ventricle (*short arrow*).

D, CT scan shows blood in the fourth ventricle (*arrow*).


The origin of intraventricular meningiomas is uncertain, although they appear to arise either from the stroma of the choroid plexus or from rests of arachnoid tissue within the choroid (4). This explains their frequent occurrence in the trigone, and at postmortem, they can be seen to arise from the choroid plexus via a vascular pedicle.

The CT appearances are similar to those of meningiomas elsewhere (1, 2). In most cases, they appear as well-defined mass lesions that show marked enhancement after intravenous contrast. Calcification is seen in as many as 50%. On T1- and T2-weighted MR images, the

lesions are usually isointense to normal brain, although the signal intensity may be markedly reduced in the presence of calcification (12). Angiographic appearances have been well described (1, 6).

The clinical presentation is variable and usually occurs when the tumors are very large (1, 4). Symptoms related to obstruction of cerebrospinal fluid flow and pressure on the surrounding brain are the most common and include postural headache, nausea, vomiting, tinnitus, syncope, hemiparesis, and focal epilepsy. Presentation with subarachnoid hemorrhage has been described (9). The clinical presentation is

Fig 2. Postcontrast CT scans correspond to Figure 1A and B. Both images show significant enhancement within the tumor.



indistinguishable from aneurysmal subarachnoid hemorrhage. In the present case and in three previous reports, there was a history of similar but less severe episodes, suggesting that repeated small bleeds may be a common feature.

Intraventricular meningioma is the most common intraventricular tumor of the trigone in adults (2, 5). The differential diagnosis must include metastatic deposits in the choroid plexus, which also may give rise to hemorrhage, although it is unusual for metastases to become so large before presentation. Primary gliomas may arise in the ventricles, although they have not been associated with hemorrhage, are rarer than meningiomas, and are more commonly found in the body of the ventricle (2, 5). Other intraventricular tumors include ependymomas, which are much more common in the ventricular body, rarely calcify, have not been associated with hemorrhage, and have a prominent cystic component in the majority of cases (2). Ventricular hemorrhage is a recognized complication of choroid plexus papilloma, which may be indistinguishable from meningioma on CT. However, these tumors are rare after adolescence, and the patients age allows a clear distinction in most cases (9). Cavernous hemangiomas rarely occur in a ventricle, may present with hemorrhage, and on CT are often round and well demarcated and of slightly uneven high density in the calcification range, but show minimal or no enhancement (13, 14). Neurocytomas are uncommon intraventricular tumors

and rarely bleed (15). Hemangioblastomas are extremely rare in a supratentorial site but have been reported in the third ventricle (16).

In conclusion, we present an intraventricular hemorrhage arising from a meningioma in the trigone of the lateral ventricle. The most common entities in the differential diagnosis of hemorrhage from intraventricular tumors in this site are choroid plexus metastasis in adults and choroid plexus papillomas in children.

References

1. Mani RL, Hedgcock MW, Mass SI, Gilmor RL, Enzmann DR, Eisenberg RL. Radiographic diagnosis of meningioma of the lateral ventricle. *J Neurosurg* 1978;49:249-255
2. Morrison G, Sobel DF, Kelley WM, Norman D. Intraventricular mass lesions. *Radiology* 1984;153:435-442.
3. Silver J, Ganti R, Hilal S. Computed tomography of tumours involving the atria of the lateral ventricles. *Radiology* 1982;145:71-78.
4. Guidetti B, Delfini R, Gagliardi FM, Vagnozzi R. Meningiomas of the lateral ventricles: clinical, neuroradiologic, and surgical considerations in 19 cases. *Surg Neurol* 1985;24:364-370
5. Jelinek J, Smirniotopoulos JG, Parisi JE, Kanzer M. Lateral ventricular neoplasms of the brain: differential diagnosis based on clinical, CT, and MR findings. *AJNR Am J Neuroradiol* 1990;11:567-574
6. Fornari M, Savoiardo M, Morello G, Solero C. Meningiomas of the lateral ventricle. *J Neurosurg* 1981;54:64-74
7. Parker D, Eschelman D, Gibbens D, Rabinov J, O'Conner J. Neuroradiology case of the day. *AJR Am J Roentgenol* 1991;156:1307-1313
8. Arseni S, Ionescu S, Meretsis M. Meningiomas of the lateral ventricle. *Psychiat Neurol Neurochir* 1968;71:319-336

9. Smith V, Stein P, McCarty C. Subarachnoid haemorrhage due to lateral ventricular meningiomas. *Surg Neurol* 1975;4:241-243
10. Markwalder T, Markwalder R, Markwalder H. Meningioma of the anterior part of the third ventricle. *J Neurosurg* 1979;50:233-235
11. Kendall B, Reider-Grosswasser I, Valentine A. Diagnosis of masses presenting within the lateral ventricles on computed tomography. *Neuroradiology* 1983;25:11-22
12. De la Sayette V, Rivaton F, Chapon F, Hubert P, Ganem F, Houtteville JP. Meningioma of the third ventricle. *Neuroradiology* 1991;33:354-356
13. Chadduck W, Binet E, Farrell F, Araoz C, Reding D. Intraventricular cavernous hemangioma: report of three cases and review of the literature. *Neurosurgery* 1985;16:189-197
14. Iwasa H, Indei I, Sato F. Intraventricular cavernous hemangioma: a case report. *J Neurosurg* 1983;59:153-157
15. Smoker W, Townsend J, Reichman M. Neurocytoma accompanied by intraventricular hemorrhage: case report and literature review. *AJNR Am J Neuroradiol* 1991;12:765-770
16. Black M, Tien R, Hesselink J. Third ventricular hemangioblastoma: MR appearance. *AJNR Am J Neuroradiol* 1991;12:553