

Generic Contrast Agents

Our portfolio is growing to serve you better. Now you have a *choice*.



[VIEW CATALOG](#)

AJNR

Contrast-enhanced magnetization transfer MR of the brain: importance of precontrast images.

J R Meyer, R W Androux, N Salamon, B Rabin, C Callahan,
T B Parrish, J Prager and E J Russell

This information is current as
of May 22, 2025.

AJNR Am J Neuroradiol 1997, 18 (8) 1515-1521
<http://www.ajnr.org/content/18/8/1515>

Contrast-Enhanced Magnetization Transfer MR of the Brain: Importance of Precontrast Images

J. R. Meyer, R. W. Androux, N. Salamon, B. Rabin, C. Callahan, T. B. Parrish, J. Prager, and E. J. Russell

PURPOSE: To determine the importance of obtaining precontrast T1-weighted magnetization transfer (MT) MR images for better interpretation contrast-enhanced T1-weighted MT images. **METHODS:** One hundred fifty-five patients referred for MR imaging of the brain were examined prospectively with noncontrast T1-weighted imaging, noncontrast T1-weighted imaging with MT, contrast-enhanced T1-weighted imaging, and contrast-enhanced T1-weighted imaging with MT. In the patients who had abnormally increased signal intensity on postcontrast images (with or without MT), the four imaging sequences were evaluated with regard to number of lesions and lesional signal intensity. For each of the sequences, two experienced neuroradiologists subjectively graded the lesions on a scale of 1 to 4 (4 being the most conspicuous) with regard to abnormally increased signal intensity. **RESULTS:** Twenty-two of the 155 patients had increased signal intensity on one or more of the postcontrast sequences. Eight of these 22 patients had increased signal intensity of one or more lesions on images without MT. All these lesions were seen better on images obtained with MT. An additional six of the 22 patients had increased signal intensity of one or more lesions on images obtained with MT that was not detected on images obtained without MT. Eight of the 22 patients had no high signal intensity on noncontrast images with or without MT. One of the eight had increased number and conspicuity of lesions on postcontrast MT images. **CONCLUSION:** A significant number of patients had increased signal intensity on noncontrast T1-weighted images with MT that was not seen on noncontrast T1-weighted images without MT. This high signal intensity was also visible on postcontrast MT images, and would have been mistaken for pathologic enhancement if noncontrast MT images had not been available for comparison.

Index terms: Brain, magnetic resonance; Magnetic resonance, magnetization transfer

AJNR Am J Neuroradiol 18:1515-1521, September 1997

Magnetization transfer (MT) is an increasingly popular magnetic resonance (MR) imaging technique that produces patterns of tissue contrast that differ substantially from those obtained with conventional spin-echo imaging. Several investigators have suggested that MT combined with paramagnetic contrast agents improves the detection of enhancing lesions, such as intracranial tumors, infection, and infarction (1-6). More specifically, several investigators have documented significant increases in contrast-to-noise ratios of

contrast-enhanced T1-weighted images with MT as compared with those of contrast-enhanced T1-weighted images without MT (1, 5, 6). Recently, others have observed significantly increased signal intensity on noncontrast T1-weighted images with MT in tuberous sclerosis and pediatric brain tumors (7, 8). This effect has the potential to influence the interpretation of contrast-enhanced MT images, because high signal intensity due to MT could be misinterpreted as pathologic contrast enhancement. The purpose of this study was to determine the importance of obtaining noncontrast T1-weighted MT images of the brain for the interpretation of contrast-enhanced T1-weighted MT images.

Patients and Methods

One hundred fifty-five patients referred for contrast-enhanced imaging of the brain were examined prospec-

Received November 27, 1996; accepted after revision March 21, 1997.

From the Departments of Radiology (J.R.M., R.W.A., N.S., B.R., C.C., T.B.P., J.P., E.J.R.) and Neurology (J.R.M.), Northwestern University Medical School, Chicago, Ill.

Address reprint requests to Joel R. Meyer, MD, Department of Radiology, Evanston Hospital, 2650 Ridge Ave, Evanston, IL 60201.

AJNR 18:1515-1521, Sep 1997 0195-6108/97/1808-1515

© American Society of Neuroradiology




Fig 1. A 30-year-old man with history of treated CNS lymphoma.

A, Noncontrast T1-weighted image shows low signal abnormality in the head of the caudate nucleus on the left. Note subtle increased signal along periphery of the lesion. There is no evidence of mass effect along the frontal horn of the left lateral ventricle.

B, Contrast-enhanced T1-weighted image shows no evidence of abnormal contrast enhancement.

C, Noncontrast T1-weighted image with MT shows increased signal in the head of the caudate nucleus accounting for the signal abnormality on the contrast-enhanced MT image (D).

D, Contrast-enhanced T1-weighted image with MT shows increased signal intensity of the head of the caudate nucleus, which might suggest abnormal contrast enhancement if noncontrast MT image (C) were not available for comparison. Image at 6 month follow-up (not shown) showed new areas of lymphoma in the left occipital lobe without recurrence in the head of the caudate nucleus.

other hand, the mobile hydrogen pool has a very narrow line width (longer T2 signal), making it possible to image.

Conventional imaging sequences are capable of showing mobile hydrogen directly. MT sequences, on the other hand, image the restricted hydrogen pool indirectly. Saturation of the bound hydrogen pool with MT pulses results in decreased signal intensity from the free hydrogen pool by the exchange of longitudinal magnetization between the two groups (9–11). The net effect is greater suppression of signal from normal brain parenchyma (more bound protons) than from pathologic brain lesions (more free protons), which may result in increased lesion conspicuity.

The level of MT contrast depends on how it is applied, the power level used, and the off-resonance frequency response. MT contrast is generated by using either off- or on-resonance pulses. In our study, these methods generated different MT contrast ratios, as implemented on each scanner.

The off-resonance pulse selectively affects the restricted hydrogen pool. The pulse is applied far enough off resonance (>1 kHz) to achieve selective saturation yet near enough to resonance to reduce the amount of power required to saturate the bound pool (9). An advantage of this method is that it is simple to implement and generally more robust. Potential drawbacks of this technique include the long

Fig 2. A 35-year-old woman with history of resection of low-grade glioma.

A, Noncontrast T1-weighted image shows a postoperative defect and signal abnormality of the left posterior frontal lobe.

B, Contrast-enhanced T1-weighted image shows no evidence of abnormal contrast enhancement.

C, Noncontrast T1-weighted image with MT shows the same area of increased signal intensity as on the contrast-enhanced MT image (D).

D, Contrast-enhanced T1-weighted image with MT shows a distinct area of increased signal intensity along the anterior margin of the resection site, which might suggest the possibility of abnormal contrast enhancement if noncontrast MT images were not obtained.





Fig 3. Mean signal intensity scores for contrast-enhanced (C) T1-weighted MR images with and without MT in eight patients who had increased signal intensity of one or more lesions on non-contrast images without MT.

duration of the pulse and the high power required to obtain proper contrast.

The on-resonance approach uses a series of nonselective RF pulses with pulse amplitudes described by the binomial expansion to produce the desired effect (12, 13). The simplest version of these jump-and-return pulses is the 1 – 1 pulse ($90_x - \tau - 90_{-x}$). The first pulse rotates all magnetization into the transverse plane. The delay time, τ , between the pulses is short, several hundred microseconds, but due to the extremely short T2 of the bound protons they dephase immediately relative to the free protons. The “return” RF pulse rotates the free protons back to the longitudinal axis, ready to be




Fig 4. A 53-year-old woman with long-standing history of multiple sclerosis. A, Noncontrast T1-weighted image shows several low signal abnormalities in the centrum semiovale, right greater than left. Note subtle increased signal in the periphery of two of these lesions on the right.

B, Contrast-enhanced T1-weighted image does not show definite abnormal contrast enhancement of the lesions.

C, Noncontrast T1-weighted image with MT shows hyperintense lesions of the periventricular white matter, accounting for the signal changes noted on the post-contrast T1-weighted image with MT (D).

D, Contrast-enhanced T1-weighted image with MT shows prominent hyperintense periventricular lesions that correspond well to abnormalities shown on proton density-weighted images. Without noncontrast MT images, it would be unclear whether these lesions represented abnormal contrast enhancement.

imaged. The bound proton signal is sufficiently decreased owing to the dephasing that occurs during the mixing period. More complicated pulse schemes can be used for optimal saturation effects. The advantage of the on-resonance pulse is that it is considerably shorter in duration than the off-resonance method and may be more energy efficient. However, the sensitivity to field homogeneity and chemical-shift differences may result in direct saturation of the fat or water peaks. The improved efficiency of the on-resonance pulse may account for the increase in MT contrast ratios observed for scanner 1.

Clinically, MT sequences have been shown to improve small-vessel conspicuity on time-of-




Fig 5. Mean signal intensity scores for contrast-enhanced (C) T1-weighted images with and without MT in six patients who had increased signal intensity of one or more lesions on noncontrast images with MT that was not detected on images without MT.




Fig 6. A 26-year-old woman with cysticercosis.

A, Contrast-enhanced T1-weighted image shows enhancing lesions in the right occipital lobe and superficial to the left insula. Possible areas of enhancement are noted in the left basal ganglia and left occipital lobe. A low-signal lesion is noted along the right frontal operculum.

B, Noncontrast T1-weighted image with MT shows areas of low signal intensity in the left basal ganglia, right frontal lobe, and right frontal operculum.

C, Contrast-enhanced T1-weighted image with MT shows several additional ring-enhancing lesions. Note the increased conspicuity of the occipital lesions compared with postcontrast T1-weighted image without MT.

flight intracranial MR angiograms; improve contrast on gradient-echo images of the cervical spine; provide tissue characterization in head and neck cancer, meningiomas, intracranial hemorrhage, wallerian degeneration in the feline visual system, and multiple sclerosis lesions; and improve detection of pathologic contrast enhancement on MR images (14–18).

It is well documented that MT sequences improve the conspicuity of contrast enhancement by suppressing background tissue signal intensity. This effect has been shown to be both dose related and synergistic (5, 19). Finelli et al (20) reported similar relative contrast improvement with triple-dose contrast-enhanced non-MT images compared with single-dose contrast-enhanced images with MT suppression. The synergistic effect is due to the fact that paramagnetic contrast agents decrease the MT effect within enhancing lesions, thereby further accentuating differences between enhancing lesions and background tissues (19). Therefore, the signal intensity of enhancing lesions is only minimally decreased with MT, much less than the background brain, which experiences a more pronounced reduction in signal intensity.

MT sequences are also highly sensitive to

differential MT ratios in normal brain and in nonenhancing pathologic brain lesions. Differential MT ratios in normal brain account for the typical gray/white matter reversal and high signal intensity of the basal ganglia observed on noncontrast T1-weighted images with MT (21). Similarly, differential MT ratios of pathologic brain lesions have recently been shown to be highly sensitive for detection of tuberous sclerosis and some brain tumors on noncontrast T1-weighted images with MT (7, 8).

A recent study by Mehta et al (6) suggested that MT pulses had little effect on noncontrast T1-weighted images in the first 14 patients of their 86-patient series. For time and cost considerations, they eliminated the noncontrast MT images in the remainder of their series. Our results differ from theirs in that a significant number of our patients (six of 22) had increased signal intensity on images obtained with MT that were not present on images obtained without MT, which influenced the qualitative interpretation of the contrast-enhanced MT images. Furthermore, eight of 22 patients had more pronounced increased signal intensity on the MT images than on the non-MT images, and this also influenced the qualitative interpretation of the contrast-enhanced MT images. This in-

creased signal intensity was related solely to the MT effect of nonenhancing tissue.

High signal on noncontrast images due to the MT effect on pathologic tissues is potentially problematic when interpreting contrast-enhanced images for the presence of abnormal enhancement. In our first group, in which the mean lesional signal intensity scores on unenhanced MT images were higher than on unenhanced non-MT images, a comparison of enhanced with unenhanced non-MT images is misleading. Simply documenting higher signal intensity on the contrast-enhanced MT images did not necessarily correlate with contrast enhancement. Rather, the mean signal intensity ratings show that a significant percentage of the signal change was due to MT effect alone.

The noncontrast MT effect was potentially even more problematic in the second group (six of 22 patients), in which high signal intensity on noncontrast images was seen only on those obtained with MT. Without a noncontrast MT image, all areas of increased signal intensity on contrast-enhanced MT images could be incorrectly attributed to enhancement, which could result in misdiagnosis of enhancement, as seen in three of six cases in which contrast-enhanced MT images showed increased signal intensity without high signal intensity on contrast-enhanced non-MT images. This might adversely affect diagnosis and treatment planning in a variety of pathologic conditions, including multiple sclerosis, brain infarction, and tumors.

Our study demonstrates the importance of comparing noncontrast with contrast-enhanced MT images. A significant number of patients had increased signal intensity on unenhanced MT images that was not identified on images obtained without MT. This increased signal contributes to the signal intensity on contrast-enhanced MT images, and could have been mistaken for enhancement if noncontrast MT images had not been available for comparison. Caution should be exercised when comparing postcontrast T1-weighted images without MT with postcontrast T1-weighted images with MT because of the potential for misdiagnosis resulting from false identification of contrast enhancement.

References

1. Finelli DA, Hurst GC, Gullapali RP, et al. Improved contrast of enhancing brain lesions on postgadolinium, T1-weighted spin-echo images with use of magnetization transfer. *Radiology* 1994; 190:553-559
2. Kurki T, Niemi P, Valtonen S. MR of intracranial tumors: combined use of gadolinium and magnetization transfer. *AJNR Am J Neuroradiol* 1994;15:1727-1736
3. Runge VM, Wells JW, Kirsch JE. Magnetization transfer and high-dose contrast in early brain infection on magnetic resonance. *Invest Radiol* 1995;30:135-143
4. Runge VM, Wells JW, Williams NM, et al. Detectability of early brain meningitis with magnetic resonance imaging. *Invest Radiol* 1995;30:484-495
5. Mathews V, King James C, Elster A, et al: Cerebral infarction: effects of dose and magnetization transfer saturation at gadolinium-enhanced MT imaging. *Radiology* 1994;190:547-552
6. Mehta R, Pike G, Haros S, et al. Central nervous system tumor, infection, and infarction: detection with gadolinium-enhanced magnetization transfer MR imaging. *Radiology*, 1995;195:41-46
7. Zimmerman R, Haselgrove J, Bilaniuk L, et al. Magnetization transfer suppression T1 weighted images without gadolinium enhancement in the diagnosis of pediatric brain tumors. *Radiology* 1995;197(P):307
8. Zimmerman RA, Girard N. Magnetization transfer in the evaluation of patients with tuberous sclerosis. *Radiology* 1995;197(P): 306
9. Wolff S, Balaban R. Magnetization transfer contrast (MTC) in magnetic resonance imaging. *Magn Reson Med* 1989;10:135-144
10. Grossman RI, Gomori JM, Ramer KN, et al. Magnetization transfer: theory and clinical applications in neuroradiology. *Radio-graphics* 1994;14:279-290
11. Balaban RS, Ceckler TL. Magnetization transfer contrast in magnetic resonance imaging. *Magn Reson Q* 1992;8:116-137
12. Yeung HN AA. Magnetization transfer contrast with periodic pulsed saturation. *Radiology* 1992;183:209-214
13. Hu BS CS, Wright GA, Nishimura DG, Macovski A. Pulsed saturation transfer contrast. *Magn Reson Med* 1992;26:231-240
14. Edelman RR, Ahn SS, Chien D, et al. Improved time-of-flight MR angiography of the brain with magnetization transfer contrast. *Radiology* 1992;184:395-399
15. Lin W, Tkach JA, Haacke EM, et al. Intracranial MR angiography: application of magnetization transfer contrast and fat saturation to short gradient-echo, velocity-compensated sequences. *Radiology* 1993;186:753-761
16. Lexa F, Grossman R, Rosenquist A. MR of wallerian degeneration in the feline visual system: characterization by magnetization transfer rate with histopathologic correlation. *AJNR Am J Neuroradiol* 1994;15:201-212
17. Loevner LA, Grossman RI, Cohen JA, et al. Microscopic disease in normal-appearing white matter on conventional MR images in patients with multiple sclerosis: assessment with magnetization-transfer measurements. *Radiology* 1995;196:511-515
18. Grossman RI. Magnetization transfer in multiple sclerosis. *Ann Neurol* 1994;36:S97-S99
19. Tantt JI, Sepponen RE, Lipton MJ, et al. Synergistic enhancement of MRI with Gd-DTPA and magnetization transfer. *J Comput Assist Tomogr* 1992;16:19-24
20. Finelli D, Hurst G, Karaman B, et al. Use of magnetization transfer for improved contrast on gradient-echo MR images of the cervical spine. *Radiology* 1994;193:163-171
21. Elster A, King J, Mathews V, et al. Cranial tissues: appearance at gadolinium-enhanced and nonenhanced MR imaging with magnetization transfer contrast. *Radiology* 1994;190:541-546