

Get Clarity On Generics

Cost-Effective CT & MRI Contrast Agents





Reply:

G. Zuccoli

AJNR Am J Neuroradiol 2011, 32 (11) E210 doi: https://doi.org/10.3174/ajnr.A2925 http://www.ajnr.org/content/32/11/E210

This information is current as of August 7, 2025.

Reply:

We thank Borges et al for their interest in our article. We agree that fornical involvement represents 1 of the unusual findings in Wernicke encephalopathy (WE).

The involvement of other anatomic substrates of memory like the thalami and the mamillary bodies represent typical MR imaging findings of WE, being present in 80% and 45% of patients, respectively. In our article, we did not find involvement of the fornix in any of the 56 patients. In our review of the literature (written before our article¹), we found only 2 of 53 patients showing fornical involvement. WE is an acute neurologic disorder characterized by changes in consciousness, ocular dysfunction, and gait disturbances. The memory deficits are characteristics of Wernicke-Korsakoff syndrome (WKS), which, in most cases, is the inevitable outcome of WE. The fornix projects mainly to the mamillary body and also to the septal region. Thus, the fornix is likely linked, to some extent, to the memory im-

pairment observed in WKS. However, on the basis of the higher incidence of signal-intensity alterations in the thalami and mamillary bodies in patients with WE, we believe that these regions are responsible for most of the memory deficits of WKS.

References

- Zuccoli G, Santa Cruz D, Bertolini M, et al. MR imaging findings in 56 patients with Wernicke encephalopathy: nonalcoholics may differ from alcoholics. AJNR Am J Neuroradiol 2009;30:171–76
- Zuccoli G, Pipitone N. Neuroimaging findings in acute Wernicke's encephalopathy: review of the literature. AJR Am J Roentgenol 2009;192:501-08

G. Zuccoli Radiology Department Children's Hospital of Pittsburgh Pittsburgh, Pennsylvania

http://dx.doi.org/10.3174/ajnr.A2925