## **Generic Contrast Agents**



Our portfolio is growing to serve you better. Now you have a *choice*.



## **Optic Canal Narrowing in Hunter Syndrome** (Mucopolysaccharidosis II)

A. Sen

AJNR Am J Neuroradiol 2013, 34 (3) E34-E35 doi: https://doi.org/10.3174/ajnr.A3499 http://www.ajnr.org/content/34/3/E34

This information is current as of May 11, 2025.

## Optic Canal Narrowing in Hunter Syndrome (Mucopolysaccharidosis II)

read with interest the article "Brain and Spinal MR Imaging Findings in Mucopolysaccharidoses: A Review<sup>»1</sup> and would like to illustrate optic canal narrowing in a patient with Hunter syndrome (mucopolysaccharidosis [MPS]-II).

A 6-year-old boy presented with significant loss of vision. He had been evaluated (at 3 years of age) for mucopolysaccharidosis; clinical phenotype, heparin sulfate in urine electrophoresis, and plasma enzyme levels (iduronate 2-sulfatase level: 4.2 nmol/h by fluorimetric assay by using 4-methyl umbelliferone; normal range: 167–475 nmol/h; control: 194.3) were suggestive of Hunter syndrome.

MR imaging of the brain showed periventricular white matter hyperintensities (Fig 1*A*–*C*), enlarged Virchow-Robin spaces (Fig 1*A*, -*B*), ventriculomegaly, and atrophy (Fig 1*A*–*C*, -*E*, -*F*), known features of mucopolysaccharidosis. The bilateral optic canals were symmetrically narrowed (Fig 1*D*–*F*), and hyperintensities were noted around both optic nerves (Fig 1*C*–*F*).

Hunter syndrome, or mucopolysaccharidosis II, is an X-linked progressive lysosomal storage disease due to deficiency of the lysosomal enzyme iduronate-2-sulfatase. Known ocular manifestations of Hunter syndrome include exophthalmos, hypertelorism, pigmentary retinopathy, uveal effusion, optic nerve

http://dx.doi.org/10.3174/ajnr.A3499

head swelling (disc edema,<sup>2</sup> possibly due to compression by scleral deposits), and optic atrophy.

Optic canal narrowing is a known feature of osteopetrosis,<sup>3</sup> and fibrous dysplasia and has been described in familial hypophosphatemic rickets, carbonic anhydrase type 2 deficiency (which includes osteopetrosis), and MPS-4.<sup>4</sup> It has not been previously reported in Hunter syndrome. Perhaps optic canal narrowing is underdiagnosed in Hunter syndrome because intellectually impaired children may not be able to describe visual loss.

## REFERENCES

- Zafeiriou DI, Batzios SP. Brain and spinal MR imaging findings in mucopolysaccharidoses: a review. AJNR Am J Neuroradiol 2013;34: 5–13
- Beck M, Cole G. Disc oedema in association with Hunter's syndrome: ocular histopathological findings. Br J Ophthalmol 1984;68:590-94
- Curé JK, Key LL, Goltra DD, et al. Cranial MR imaging of osteopetrosis. AJNR Am J Neuroradiol 2000;21:1110–15
- Vougioukas VI, Berlis A, Kopp MV, et al. Neurosurgical interventions in children with Maroteaux-Lamy syndrome: case report and review of the literature. *Pediatr Neurosurg* 2001;35:35–38

**A. Sen** Radiodiagnosis Devi Scans Kerala, India



**FIG 1.** Axial T2 (A) and axial T2 FLAIR (B) images show periventricular white matter hyperintensities (*arrows*), prominent Virchow-Robin spaces (*arrowheads*), ventriculomegaly, and brain atrophy. C, Coronal T2 image shows hyperintensities in periventricular white matter (*arrows*) and perioptic spaces (*arrowheads*), ventriculomegaly, and brain atrophy. Axial T2 (D) and right (E) and left (F) sagittal T2 images show narrowing of the optic canals (*arrows*), hyperintensities in perioptic spaces, and brain atrophy.