



Get Clarity On Generics

Cost-Effective CT & MRI Contrast Agents



FRESENIUS
KABI

WATCH VIDEO

AJNR

Asymptomatic Interhypothalamic Adhesions in Children

M.T. Whitehead and G. Vezina

AJNR Am J Neuroradiol 2016, 37 (4) E35

doi: <https://doi.org/10.3174/ajnr.A4703>

<http://www.ajnr.org/content/37/4/E35>

This information is current as
of August 12, 2025.

Asymptomatic Interhypothalamic Adhesions in Children

We have several comments regarding the article “Asymptomatic Interhypothalamic Adhesions in Children.”¹ We agree with the main message of the article: Referable hypothalamic–pituitary axis symptoms are rare in patients with interhypothalamic adhesions. However, because symptoms can be present on occasion, it is prudent to exclude endocrinopathy on clinical grounds. We have encountered a few cases of patients with interhypothalamic adhesions and pituitary axis disturbances, one associated with Kallmann syndrome²; 2 with septo-optic dysplasia; and 1, with abnormal weight gain.

We agree with the authors’ theory that interhypothalamic adhesions may be the result of “incomplete hypothalamic cleavage, failed apoptosis, or abnormal neuronal migration” and acknowledge the association with “gray matter heterotopia.” Therefore, additional midline abnormalities would be expected. However, the authors did not identify additional abnormalities in most patients. Nonetheless, concurrent gray matter heterotopia was present in 40%, a considerably large percentage of patients, and they proposed that heterotopia associated with interhypothalamic adhesions may be part of an unknown genetic disorder.


All portions of the brain, including the midline, must be carefully examined in patients with interhypothalamic adhesions because they represent a potential marker for brain malformation. Additional midline anomalies/abnormalities are quite common in our experience.^{2–4} These may be subtle and insignificant (hypoplasia of the falx, underrotated hippocampi, and so forth) or obvious and potentially of great consequence (malformations of


brain development).^{2–4} Indeed, review of Fig 1 demonstrates subtle midline anomalies not mentioned in the article, including hypogenesis or volume loss of the splenium (Figs 1A and E) and a partially fenestrated, persistent cavum septum pellucidum (Fig 1D).¹ In normal brains, the callosal splenium is typically equal to or larger in caliber than the genu. While a cavum septum pellucidum is a normal variation, it is uncommon in the general population beyond the neonatal period.

We strongly believe that the midline should be closely scrutinized for additional anomalies/abnormalities in patients with an interhypothalamic adhesion. Only after the brain has been carefully examined and signs/symptoms have been carefully considered can an interhypothalamic adhesion be considered an incidental and isolated finding.

REFERENCES

1. Ahmed FN, Stence NV, Mirsky DM. **Asymptomatic interhypothalamic adhesions in children.** *AJNR Am J Neuroradiol* 2015 Dec 3. [Epub ahead of print] [CrossRef Medline](#)
2. Whitehead MT, Angel JD. **Interhypothalamic adhesion in a 9-month-old male with cleft palate.** *Case Rep Radiol* 2013;2013:197415 [CrossRef Medline](#)
3. Whitehead MT, Vezina G. **Interhypothalamic adhesion: a series of 13 cases.** *AJNR Am J Neuroradiol* 2014;35:2002–06 [CrossRef Medline](#)
4. Whitehead MT, Lee B. **Neuroimaging features of San Luis Valley syndrome.** *Case Rep Radiol* 2015;2015:748413 [CrossRef Medline](#)

 **M.T. Whitehead**

 **G. Vezina**

Department of Neuroradiology
Children’s National Medical Center
Washington, DC

<http://dx.doi.org/10.3174/ajnr.A4703>