



Discover Generics

Cost-Effective CT & MRI Contrast Agents



WATCH VIDEO

AJNR

Reply:

N. Larsen, O. Jansen, C.H. Riedel, C. von der Brelie and C. Flüh

AJNR Am J Neuroradiol 2018, 39 (12) E131

doi: <https://doi.org/10.3174/ajnr.A5860>

<http://www.ajnr.org/content/39/12/E131>

This information is current as
of June 5, 2025.

REPLY:

We thank Dr Corrêa and colleagues for their interest in providing feedback for our recent article, “Vessel Wall Enhancement in Unruptured Intracranial Aneurysms: An Indicator for Higher Risk of Rupture? High-Resolution MR Imaging and Correlated Histologic Findings.”¹ We described our findings in MR vessel wall imaging in unruptured middle cerebral artery aneurysms and correlated histologic features and proposed a possible association among aneurysm wall enhancement attributable to inflammatory cell invasion, wall degeneration, and higher risk of rupture.

Dr Corrêa and colleagues pointed out that mural enhancement might not be associated with a higher risk of progression and rupture in endovascularly treated aneurysms.

We agree that wall enhancement in intracranial vessels is not necessarily pathologic or associated with a poorer prognosis. Endovascular treatment of aneurysms with coiling or flow diversion substantially alters hemodynamic, morphologic, and subsequently histologic conditions in the aneurysm wall and the adjacent segments of the parent vessel. Mural enhancement in these patients might still be of diagnostic value, but the lack of long-term follow-up data addressing this issue makes it difficult, for the time being, to assess its relevance.

Moreover, mural enhancement occurs in intracranial vessels in proximity to the dural penetration and is believed to be attributable to the presence of the vasa vasorum.² Therefore, the significance of wall enhancement in unruptured aneurysms located in these vessel segments (eg, paraophthalmic internal carotid artery, posterior inferior cerebellar artery origin) ought to be critically and separately investigated.

Furthermore, among intracranial aneurysms, various entities have been described that possibly differ in pathogenesis, morpho-

logic features, progression rate, and clinical course.³⁻⁵ In our opinion, further investigation of distinct MR vessel wall imaging features in saccular-versus-fusiform, sidewall-versus-bifurcation, and small-versus-large aneurysms is in order.

REFERENCES

1. Larsen N, von der Brelie C, Trick D, et al. **Vessel wall enhancement in unruptured intracranial aneurysms: an indicator for higher risk of rupture? High-resolution MR imaging and correlated histologic findings.** *AJNR Am J Neuroradiol* 2018;39:1617–21 [CrossRef Medline](#)
2. Portanova A, Hakakian N, Mikulis DJ, et al. **Intracranial vasa vasorum: insights and implications for imaging.** *Radiology* 2013;267:667–79 [CrossRef Medline](#)
3. Baharoglu MI, Lauric A, Gao BL, et al. **Identification of a dichotomy in morphological predictors of rupture status between sidewall- and bifurcation-type intracranial aneurysms.** *J Neurosurg* 2012;116:871–81 [CrossRef Medline](#)
4. Wiebers DO, Whisnant JP, Huston J 3rd, et al. International Study of Unruptured Intracranial Aneurysms Investigators. **Unruptured intracranial aneurysms: natural history, clinical outcome, and risks of surgical and endovascular treatment.** *Lancet* 2003;362:103–10 [Medline](#)
5. Varble N, Tutino VM, Yu J, et al. **Shared and distinct rupture discriminants of small and large intracranial aneurysms.** *Stroke* 2018;49:856–64 [CrossRef Medline](#)

 **N. Larsen**

 **O. Jansen**

 **C.H. Riedel**

Department of Radiology and Neuroradiology
University Hospital Schleswig-Holstein Campus Kiel
Kiel, Germany

 **C. von der Brelie**

Department of Neurosurgery
University Hospital Göttingen
Göttingen, Germany

 **C. Flüh**

Department of Neurosurgery
University Hospital Schleswig-Holstein Campus Kiel
Kiel, Germany

<http://dx.doi.org/10.3174/ajnr.A5860>