



Discover Generics

Cost-Effective CT & MRI Contrast Agents



FRESENIUS
KABI

WATCH VIDEO

AJNR

Cerebral varix and probable venous angioma: an unusual isolated anomaly.

J D Meyer, P Baghai and R E Latchaw

AJNR Am J Neuroradiol 1983, 4 (1) 85-87

<http://www.ajnr.org/content/4/1/85.citation>

This information is current as
of June 20, 2025.

Cerebral Varix and Probable Venous Angioma: An Unusual Isolated Anomaly

J. D. Meyer,¹ P. Baghai,² and R. E. Latchaw¹

A patient with headaches was found at angiography to harbor a venous saclike aneurysm unassociated with any evidence of arteriovenous shunting. Computed tomographic (CT) and angiographic findings are described. No previous report with similar radiographic findings has been found.

Case Report

A 31-year-old man came to the Veterans Administration Medical Center after 3–4 years of right frontal headaches that had been increasing in frequency and intensity. No other complaints were noted. There was no history of migraine and family history was unremarkable. Physical examination including neurologic testing was completely normal. Initial CT examination (not shown) was suspicious for an intracranial aneurysm, and an arteriogram was obtained (fig. 1). The arterial phase on the right side was normal in appearance. Appearing late in the venous phase was a smooth, round, saclike vein that drained a portion of the caudate nucleus and lateral basal ganglia on the right side, and emptied by way of a large single vessel into the sylvian venous system. The appearance of several of the small veins draining the lateral basal ganglia and emptying into the sac was noted. A lateral scout film after the anteroposterior film sequence (the examination was partly performed in a single plane system) revealed persistent opacification of this saclike vessel for several minutes. Subsequent higher resolution CT examination (fig. 2) on the General Electric 8800 scanner revealed a rounded lesion of minimal hyperdensity on the noncontrast portion that enhanced considerably after contrast administration. The two major vessels draining into this venous sac were also noted and corresponded to angiographic changes. In addition, the caudate nucleus and a part of the lateral basal ganglia appeared to enhance somewhat more than the opposite side. The examination revealed on both sides and in the posterior circulation no evidence of arteriovenous malformation or any other vascular abnormality. There was no sign of vasculitis. The patient was treated conservatively and no operation was performed.

Discussion

Nonneoplastic vascular malformations can be generally classified into five categories as described by McCormick

[1]. This classification includes telangiectasias, varix, cavernous malformation or angioma, arteriovenous malformation, and venous malformation. The classification used by Russell and Rubinstein [2], summarized by Wendling et al. [3], and cited by others [4–8], deletes the category of varix. In a comprehensive review by Noran [9], cerebral varix is discussed at length with descriptions of cases reported much earlier. Included in these descriptions are those composed of lesions containing more than one abnormal vein. These might possibly have represented arteriovenous malformations. However, there are well documented cases in which a single large vessel was encountered as an isolated phenomenon; that is, one not associated with arteriovenous malformation at necropsy [10–13]. These varices demonstrated a tendency to rupture, but the true likelihood of rupture was not known. Noran reported no specific site of predilection and a tendency for association with a dural sinus. Histologically, a relatively large, thin-walled vessel was found, usually lined by a single layer of endothelium and circled by a single layer of thin connective tissue. Muscle fibers and elastic tissue were generally absent.

Arteriographic examination on these cases from the earlier literature and from the three cases added by Noran is unavailable. Cushing and Bailey [14] clearly described a large, isolated varix and categorically separated it from lesions composed of multiple veins, arteries and veins, or telangiectasis.

The etiology of the varix has been only speculative and authors appear to be unsure in even classifying the abnormality. It appears to be a rather uncommon entity. No association with the more typical venous angioma, now well described in the literature [4–8, 15–27], has been noted. Our case appears to have some characteristics suggesting that it may represent a variation of venous angioma. The caudate nucleus and lateral basal ganglia are slightly hypervascular, and this hypervascularity in turn is related to prominent veins seen both on angiography and CT. These veins drain into the varix, which then empties into the sylvian group of vessels, part of the superficial venous system. One

Received March 1, 1982; accepted after revision May 28, 1982.

¹Department of Radiology, Division of Neuroradiology, University of Pittsburgh School of Medicine, Presbyterian-University Hospital, DeSoto at O'Hara Sts., Pittsburgh, PA 15213. Address reprint requests to J. D. Meyer.

²Department of Neurological Surgery, University of Pittsburgh School of Medicine, Montefiore Hospital, Pittsburgh, PA 15213.




Fig. 1.—Anteroposterior (A) and lateral (B) angiograms. Venous phase of right-sided injection reveals rounded saclike venous aneurysm (arrows) and drainage by way of large single vein (arrowheads) into sylvian venous system. C, Slightly oblique and enlarged. Several prominent small vessels (arrowheads) draining into venous aneurysm are suggestive of venous angioma. D, Lateral scout film before injection of opposite carotid. Persistent contrast in lower part of venous aneurysm has a prominent semilunar shape.




Fig. 2.—A, Noncontrast scan through venous aneurysm (arrowhead). B, After administration of contrast material. Dense enhancement of aneurysm and visualization of two major veins entering it (arrow). C, Just above venous sac. Somewhat increased enhancement in right caudate nucleus and in lateral basal ganglia, excluding thalamus (arrow).

would expect the basal ganglia drainage to be into the deep system. On the angiogram several prominent veins appear to be related to the varix, but their appearance is not wholly typical of venous angioma: they are relatively fewer in number and do not present a collection of parallel channels, the "spoke-wheel" appearance often seen. The appearance also is not that of the cavernous angioma, which may demonstrate calcification, mass effect, occasional staining, and vascular pooling of contrast material, with possible dense enhancement on CT [28-37].

The findings in our case appear to clearly represent a cerebral varix. Association with a venous angioma has not been previously reported or noted by earlier authors. The evidence contained herein would suggest that the association of venous angioma and varix may be present.

Varices have been noted often with high-flow vascular malformations. They are the most prominent component of the vein of Galen fistula seen in children and occasionally adults and are noted uncommonly in the orbit.

REFERENCES

- McCormick WF. The pathology of vascular ("arteriovenous") malformations. *J Neurosurg* 1966;24:807-816
- Russell DS, Rubinstein LJ. *Pathology of tumors of the nervous system*, 3d ed. Baltimore: Williams & Wilkins, 1971:85-108
- Wendling LR, Moore JS, Kieffer SA, Goldberg HI, Latchaw RE. Intracerebral venous angioma. *Radiology* 1976;119:141-147
- Michels LG, Bentson JR, Winter J. Computed tomography of cerebral venous angiomas. *J Comput Assist Tomogr* 1977;1:149-154
- Maehara T, Tasaka A. Cerebral venous angioma: computerized tomography and angiographic diagnosis. *Neuroradiology* 1978;16:296-298
- Fierstein SB, Pribram HW, Hieshima G. Angiography and computed tomography in the evaluation of cerebral venous malformations. *Neuroradiology* 1979;17:137-148
- Pertain CL, Guinto FC, Scatliff JH, Limbacher J, Janicki T, Heindel CC. Cerebral venous angioma: correlation of radionuclide brain scan, transmission computed tomography, and angiography. *J Nucl Med* 1979;20:1166-1169
- Moritake K, Handa H, Mori K, Ishikawa M, Morimoto M, Takebe Y. Venous angiomas of the brain. *Surg Neurol* 1980;14:95-105
- Noran HH. Intra-cranial vascular tumors and malformations. *Arch Pathol Lab Med* 1945;39:393-416
- Amsler C. Über intrakranielle Varizenbildung an der Vena ophthalmomeningea Hyrtl. *Z Pathol* 1912;11:254-261
- Beger H. Kasuistischer Beitrag zur cerebralen Varizenbildung. *Virchows Arch Pathol* 1921;231:439-452
- Anders O. Über einen Fall von ausgedehnter cerebraler Varizenbildung mit tödlicher Blutung in Verbindung mit Sinus pericranii. *Beitr Pathol* 1918;64:540-550
- Pfannenstiel. Apoplexie als tödlicher Ausgang von Eklampsie. *Zentralbl Gynaekol* 1887;11:601-607
- Cushing H, Bailey P. *Tumors arising from the blood vessels of the brain*. Springfield, IL: Thomas, 1928:13-16
- Scotti LN, Goldman RL, Rao GR, Heinz ER. Cerebral venous angioma. *Neuroradiology* 1975;9:125-128
- Preissig RS, Preissig SH, Goree JA. Angiographic demonstration of a cerebral venous angioma. *J Neurosurg* 1976;44:628-631
- Sarwar M, McCormick WF. Intracerebral venous angioma. *Arch Neurol* 1978;35:323-325
- Sartor VK, Flidner E, Weber K. Venose Angiome des Gehirns. *ROFO* 1978;128:171-176
- Cabanes J, Blasco R, Garcia M, Tamarit L. Cerebral venous angiomas. *Surg Neurol* 1979;11:385-389
- Iraci G, Galligioni F, Gerosa M, et al. Intracerebral venous angiomas as a cause of exophthalmos. *Ann Ophthalmol* 1979;11:603-612
- Kasamo S, Kobayashi E, Awa H, et al. A case of cerebral venous angioma as an epileptogenic lesion detected by CT scan and surgically treated. *Neurol Med Chir (Tokyo)* 1980;20:865-873
- Pardatscher K, Fiore DL, Galligioni F, Iraci G. Diagnosis of cerebral venous angioma by rapidly enhanced CT scan. *Surg Neurol* 1980;14:111-113
- Solomon EH, Bonstelle CT, Modic MT, Kaufman B. Angiographic and computed tomographic correlation in cerebral venous angiomas. *CT* 1980;4:217-221
- Tomaccini D, Venturi C, Scarfo GB. Cerebral venous angioma, cutaneous angioma, facial asymmetry and recurrent stroke. *Eur Neurol* 1981;20:29-32
- Branch CE Jr, Kunath AM, Buscemi JH. Sublingual venous angioma—marker of intracranial lesion? *Arch Neurol* 1981;38:259-260
- Hacker DA, Latchaw RE, Chou SN, Gold LHA. Bilateral cerebellar venous angioma. *J Comput Assist Tomogr* 1981;5:424-426
- Saito Y, Kobayashi N. Cerebral venous angiomas. *Radiology* 1981;139:87-94
- Jonutis AJ, Sondheimer FK, Klein HZ, Wise BL. Intracerebral cavernous hemangioma with angiographically demonstrated pathologic vasculature. *Neuroradiology* 1971;3:57-63
- Segall HD, Segal HL, Teal JS, Rumbaugh CL, Bergeron RT. Calcifying cerebral cavernous hemangioma with brain scan and angiographic findings. *Neuroradiology* 1974;7:133-138
- Bartlett JE, Kishore PRS. Intracranial cavernous angioma. *AJR* 1977;128:653-656
- Numaguchi Y, Fukui M, Miyake E, et al. Angiographic manifestations of intracerebral cavernous hemangioma. *Neuroradiology* 1977;14:113-116
- Kawai K, Fukui M, Tanaka A, Kuramoto S, Kitamura K. Extracerebral cavernous hemangioma of the middle fossa. *Surg Neurol* 1978;9:19-25
- Ramina R, Ingunza W, Vonofakos D. Cystic cerebral cavernous angioma with dense calcification. *J Neurosurg* 1980;52:259-262
- Ishikawa M, Handa H, Moritake K, Mori K, Nakano Y, Aii H. Computed tomography of cerebral cavernous hemangiomas. *J Comput Assist Tomogr* 1980;4:587-591
- Galzio RJ, Zenobii M, Carbonin G, Cristuibi-Grizzi L. Hemangioma calcificans. *Surg Neurol* 1980;14:331-335
- Pozzati E, Padovani R, Morrone B, Finizio F, Gaist C. Cerebral cavernous angiomas in children. *J Neurosurg* 1980;53:826-832
- Bogren H, Svalander C, Wickbom I. Angiography in intracranial cavernous hemangiomas. *Acta Radiol [Diagn] (Stockh)* 1970;10:81-89