Generic Contrast Agents

Our portfolio is growing to serve you better. Now you have a choice.





Reply:

T.A. Kennedy

AJNR Am J Neuroradiol 2019, 40 (7) E37 doi: https://doi.org/10.3174/ajnr.A6094 http://www.ajnr.org/content/40/7/E37

This information is current as of May 10, 2025.

REPLY:

Te appreciate the thoughtful comments shared by De Bernardo et al regarding our recent publication exploring the use of contrast-enhanced 3D-FLAIR imaging in the evaluation of patients with idiopathic intracranial hypertension (IIH).¹ In their comments, the authors of the response wished to raise awareness of how patients with IIH could be serially monitored with sonography as a more sensitive and cost-effective technique to evaluate optic nerve edema compared with contrast-enhanced MR imaging.

We would like to highlight, however, that the overarching objective of our report was to evaluate the sensitivity of contrastenhanced 3D-FLAIR imaging in patients for the initial (and sometimes opportunistic) diagnosis of IIH, as opposed to serial and longitudinal imaging of these patients. In patients presenting a priori with headaches and blurred vision, it is important to

evaluate the brain to rule out other causes of elevated intracranial pressure, and a head MR imaging without and with contrast remains the standard of care to evaluate this patient population.² Our investigation identified a moderate correlation between the Frisén Scale and findings seen on contrast-enhanced 3D-FLAIR, particularly in patients with higher grades of papilledema. Once a diagnosis of IIH has been clinically established, sonography may play a role in follow-up as the authors suggest.

REFERENCES

- 1. Golden E, Krivochenitser R, Mathews N, et al. Contrast-enhanced 3D-FLAIR imaging of the optic nerve and optic nerve head: novel neuroimaging findings of idiopathic intracranial hypertension. AJNR Am J Neuroradiol 2019;40:334-39 CrossRef Medline
- 2. American College of Radiology. ACR Appropriateness: Headache. (https://acsearch.acr.org/docs/69482/Narrative/). Accessed May 10,

T.A. Kennedy

Department of Radiology, Division of Neuroradiology University of Wisconsin School of Medicine and Public Health Madison, Wisconsin

http://dx.doi.org/10.3174/ajnr.A6094