

Generic Contrast Agents

Our portfolio is growing to serve you better. Now you have a choice.



FRESENIUS
KABI

[VIEW CATALOG](#)

AJNR

Erratum

AJNR Am J Neuroradiol 2019, 40 (7) E39

doi: <https://doi.org/10.3174/ajnr.A6082>

<http://www.ajnr.org/content/40/7/E39>

This information is current as
of May 9, 2025.

In the article “Prevalence of Superior Semicircular Canal Dehiscence on High-Resolution CT Imaging in Patients without Vestibular or Auditory Abnormalities” by A.W. Berning, K. Arani, and B.F. Branstetter, IV (*AJNR Am J Neuroradiol* 2019;40:709–12), the images and legends for Figs 3 and 4 were incorrectly matched. They should appear as shown below. The authors regret the error.

<http://dx.doi.org/10.3174/ajnr.A6082>



FIG 3. A thin-but-intact superior semicircular canal. Coronally reformatted CT of the temporal bone demonstrates very thin bone overlying the superior semicircular canal (*arrow*). This quantity of bone should be classified as intact in patients with and without suggestive symptoms.




FIG 4. Dehiscent superior semicircular canal. Coronally reformatted CT of the temporal bone demonstrates dehiscence of the roof of the superior semicircular canal (*arrow*).