

Get Clarity On Generics

Cost-Effective CT & MRI Contrast Agents



FRESENIUS
KABI

WATCH VIDEO

AJNR

**Celebrating 35 Years of the AJNR: May 1987
edition**

AJNR Am J Neuroradiol 2022, 43 (5) 789

doi: <https://doi.org/10.3174/ajnr.P6842>

<http://www.ajnr.org/content/43/5/789.citation>

This information is current as
of August 20, 2025.

Celebrating 35 Years of the AJNR

May 1987 edition


Study of Movement Disorders and Brain Iron by MR

J. Neal Rutledge^{1,2}
Sadek K. Hilal¹
A. John Silver¹
Richard Defendini³
Stanley Fahn⁴


Heavily T2-weighted high-field MR images provide a unique opportunity for the evaluation of the extrapyramidal motor system. The images are affected by the presence of small amounts of naturally occurring paramagnetic substances—principally iron—that delineate the neostriatum (caudate and putamen), globus pallidus, red nucleus, substantia nigra, and dentate nucleus, primarily by a decrease in signal secondary to the T2* effect. Movement disorders are associated with either increased or decreased signal or both in these structures, depending on the pathologic process. In the initial evaluation of 113 patients with a variety of movement disorders, good correlation of imaging abnormalities can be made with a simplified schema of the extrapyramidal pathways and a system of classification of abnormal movements, parkinsonism/tremor, dystonia, chorea, myoclonus, and hemiballismus. Parkinsonisms are characterized by abnormalities of the cortico-ponto-cerebello-dentato-rubro-thalamo-cortico-spinal tract¹ or the nigrostriatal tract. Dystonias are characterized by abnormalities of the neostriatum predominantly affecting the putamen. Chorea are also characterized by abnormalities of the neostriatum but predominantly affecting the caudate nucleus. Hemiballismus is characterized by lesions affecting the subthalamic nucleus or associated pathway.

Iron

Interest in movement disorders is heightened by the recent ability to map the distribution of macromolecular complexes of Fe(II) in the brain with heavily T2-weighted images. This is accomplished through contrast created by the T2* effect, a local inhomogeneity in the magnetic field that dephases spin and results in loss of signal [1]. This effect is different from the paramagnetic effect of smaller soluble contrast agents, such as gadolinium, and corresponds roughly with the ferric iron



D



This article appears in the May/June 1987 issue of AJNR and the August 1987 issue of AJNR. Received July 10, 1986; accepted after revision November 20, 1986.

Presented at the annual meeting of the American Society of Neuroradiology, San Diego, January 1987.

This work was supported in part by National Institutes of Health grant CA 28981 and by Philips Medical Systems, Inc.

¹Department of Radiology, Division of Neuro-radiology, Columbia-Presbyterian Medical Center, New York, NY 10032.

²Present address: 10 Medical Arts Square, Austin, TX 78705. Address reprint requests to J. N. Rutledge.

³Department of Pathology, Division of Neuro-pathology, Columbia-Presbyterian Medical Center, New York, NY 10032.

⁴Department of Neurology, Division of Movement Disorders, Columbia-Presbyterian Medical Center, New York, NY 10032.

AJNR 6:397-411, May/June 1987
0195-6108/87/0603-0397
© American Society of Neuroradiology

Radiation Effects on Cerebral White Matter: MR Evaluation


Jay S. Tsuruda^{1,2}
Keith E. Kortman^{1,3}
William G. Bradley^{1,3}
Donald C. Wheeler¹
William Van Dalsen¹
Thomas P. Bradley¹

The purpose of this study was to evaluate the white-matter changes associated with external beam radiation for a wide variety of central nervous system tumors. Moderately T2-weighted spin-echo images with a 2000-msec repeat and 56-msec-echo time were analyzed for white-matter abnormalities without knowledge of the patient's history. These were correlated with radiation dose, port, and time since completion of therapy, and then compared with an age-matched control group of 180 patients with nonradiated, space-occupying, intracranial lesions. Radiatic lesions were characterized as symmetric, high-signal foci in the periventricular matter. Relative sparing of the posterior fossa, basal ganglia, and internal capsule noted. In patients older than 20 years, these changes paralleled those seen in but were more prevalent ($p < .005$). In 25 patients with sequential MRI, scan findings remained stable. In those patients with limited treatment fields, for pituitary adenomas, no statistical differences were seen between radiation-treated and nontreated groups. Cerebral white-matter changes that mimic deep white-matter are frequently seen in response to therapeutic radiation. There is a incidence of radiation effects, becoming more marked in older patients. MR must consider the neuropathologic consequences of therapeutic radiation, white demyelination, microvascular occlusion, and blood-brain barrier breakdown.

The goal of therapeutic radiology is to provide a sufficient dose to treat nervous system (CNS) tumors without affecting adjacent healthy tissue. Therapeutic regimens have been implemented, long-term effects of neurotoxicity have been recognized [1-3]. A spectrum of CNS has been noted on CT including atrophy, decreased attenuation of the deep white matter, and rarely focal, enhancing radionecrosis [4-7]. The first neurotoxicity have been correlated with clinical findings in both patients and experiments [1]. More recently, similar findings of radiotherapy-related changes in the brain have been described with MR [8-10]. Of interest, is the formation white-matter periventricular foci characterized by prolonged T1 and T2 times compared with normal white matter [10] in a pattern suggestive of demyelination or ischemia. Both can be identified readily by MR [11]. The natural prevalence and severity of focal periventricular abnormalities (e.g., deep white-matter infarction) increase with age [11, 12]; these abnormalities must be considered before ascribing changes to radiotherapy. For this reason, a retrospective review of a large patient population was undertaken; MR was used to further characterize the effects of radiation on cerebral white matter with respect to age, prevalence, time of onset, and both the temporal and spatial stability of these lesions.

Materials and Methods

MR imaging was performed using a Diasonics MT/S system operating at 0.35 T. Multiple-



This article appears in the May/June 1987 issue of AJNR and the July 1987 issue of AJNR. Received August 20, 1986; accepted after revision December 16, 1986.

Presented at the annual meeting of the American Roentgen Ray Society, Washington, DC, April 1986.

¹NMR Imaging Laboratory, Huntington Medical Research Institutes, 10 Pho St., Pasadena, CA 91105. Address reprint requests to W. G. Bradley.

²Present address: Department of Radiology, University of California, San Francisco, CA 94143.

³Department of Radiology, Huntington Memorial Hospital, Pasadena, CA 91105.

AJNR 6:431-437, May/June 1987
0195-6108/87/0603-0431
© American Society of Neuroradiology