



Get Clarity On Generics

Cost-Effective CT & MRI Contrast Agents



FRESENIUS
KABI

WATCH VIDEO

AJNR

This information is current as
of August 22, 2025.

Reply

M. Petruzzellis, R. De Blasi, V. Lucivero, M. Sancilio, M. Prontera, A. Tinelli, D.M. Mezzapesa and F. Federico

AJNR Am J Neuroradiol published online 26 September 2007

<http://www.ajnr.org/content/early/2007/10/04/ajnr.A0731.citation>

Reply:

We thank Dr. Fox for his comments concerning our previously published letter to the editor entitled, "Cerebral Aneurysms in a Patient with Osteogenesis Imperfecta and Exon 28 Polymorphism Of COL1A2."¹ He strongly informs us of a misunderstanding regarding the de novo aneurysm that developed after 4 months. Dr. Fox points out some potential errors in our diagnosis, claiming that the de novo aneurysm is actually a refilling of the previously treated one. We believe that indeed at 4 months, our images revealed that a new aneurysm had developed in front of the previous one (Fig 2B). Further evidence of this was found in the posttreatment un-subtracted images (not published due to space constraints), in which it was possible to appreciate the stent crossing the vertebrobasilar junction and the coils occluding 2 different and clearly separable aneurysms.

Finally, we are grateful to Dr. Fox for bringing to our attention some bibliographic references that may help us achieve a better understanding of the anatomic features of this particular case.

Reference

1. Petruzzellis M, De Blasi R, Lucivero V, et al. Cerebral aneurysms in a patient with osteogenesis imperfecta and exon 28 polymorphism of COL1A2. *AJNR* Am J Neuroradiol 2007;28:397-98

M. Petruzzellis
R. De Blasi
V. Lucivero
M. Sancilio
M. Prontera
A. Tinelli

D.M. Mezzapesa
F. Federico

*Dipartimento di Scienze Neurologiche e Psichiatriche
Università degli Studi di Bari-Italia
Bari, Italy*

DOI 10.3174/ajnr.A0731