



Discover Generics

Cost-Effective CT & MRI Contrast Agents



[VIEW CATALOG](#)

AJNR

CTP: Time for Patient-Tailored Acquisition Protocols

J. Molad

AJNR Am J Neuroradiol published online 24 February 2022
<http://www.ajnr.org/content/early/2022/02/24/ajnr.A7457>

This information is current as
of September 11, 2025.

CTP: Time for Patient-Tailored Acquisition Protocols

The use of perfusion imaging has greatly increased during the recent decade, becoming an integral part of the acute stroke imaging protocol,¹ and it is now included in recent acute stroke guidelines, recommending perfusion imaging in the late (6- to 24-hour) timeframe for intervention.² Because the role of perfusion imaging in patient selection is increasingly widespread, obtaining high-quality and accurate acquisitions and analysis becomes of critical importance.

Most stroke centers worldwide use CTP rather than MR perfusion due to its wider availability, rapid imaging, superior vascular imaging of extracranial and distal intracranial vessels, and fewer absolute contraindications.¹⁻³ However, CTP still has several limitations and pitfalls, precluding it from being the criterion standard for the assessment of salvageable brain tissue. These limitations include lack of standardized acquisition protocols, heterogeneity of postprocessing techniques and thresholds, inaccurate estimation of white matter lesions, and low sensitivity for the detection of lacunar infarcts.⁴⁻⁶

Delayed arrival of the contrast bolus or truncation (early termination) is the main cause of inadequate acquisition of CTP and is reported to occur in up to 67% of CTP scans.^{7,8} Truncation may result in incomplete capture of the tissue time-attenuation curves and thus precludes accurate estimation of the infarct core and penumbra. Previous studies have found truncation to falsely repartition the ROI into a larger ischemic core and smaller penumbra volumes.⁸ Delayed contrast arrival and early termination frequently result from the susceptibility of CTP to the influence of impaired cardiac output or flow-limiting carotid artery stenosis.^{4,9} Because the 2 leading etiologies for large-vessel occlusion (LVO) are large-vessel atherosclerosis and atrial fibrillation, inadequate contrast-injection timing becomes a main pitfall of CTP. Truncation rates vary in accordance with the duration of CTP scan times. Although a longer scan duration of 90 seconds may reduce truncation rates many centers use 40- to 60-second protocols to reduce the radiation load and risk of patient movement.¹⁰

In this issue of the *American Journal of Neuroradiology*, Hartman et al¹¹ examined the use of patient-tailored CTP acquisition protocols, a standard acquisition versus specific low cardiac output (LCO) protocol, to achieve better timing of the contrast bolus and avoid truncation.¹¹ This novel approach is in contrast

to previous studies¹⁰ that tried to define a unified optimal scan duration. In the current trial, the LCO protocol had a prolonged image-acquisition time of 75 seconds and an expanded scan delay of 7 seconds after contrast bolus injection, compared with an acquisition time of 65 seconds and a delay of 2 seconds in the standard protocol. Protocol selection was determined using test dose enhancement rise timing, routinely performed on CTA. A cutoff value of 15 seconds from contrast injection to a measure of 100 HU in the aortic arch was used. Among 157 scans obtained, truncation was demonstrated in only 2.5% versus 9.8% among 153 patients who underwent routine CTP using the standard protocol exclusively. The use of this semiautomated model allowed substantial improvement in the CTP acquisition without exposing patients to unneeded excess radiation. Most important, among patients who were found suitable for the standard acquisition duration, no cases of truncation were documented. Therefore, it seems that the proposed algorithm for protocol selection has very good sensitivity and may imply that the extension of the scan duration in the LCO protocol may reduce truncation rates even further.

The authors further examined factors that distinguish patients with high-versus-low risk for early termination. Age, a low left-ventricular ejection fraction, and the absence of hypertension were found to be independent risk factors for truncation. These factors and other previously mentioned parameters are directly associated with contrast arrival time to the intracranial vessels.^{4,9,11} Such clinical characteristics may also be used to detect patients with a high risk of truncation. However, the use of test dose enhancement rise timing is preferable, due to the semiautomated nature of the described protocol, which requires no preliminary clinical data nor any action from the stroke physician and thus would not lead to any delay in imaging or intervention times.

The limitations of the current study are mainly because all scans were performed in a single center with the use of a single CTP postprocessing software package. The suggested protocols should be examined using various scanners and software programs to assess the generalizability of the results. Second, a relatively high exclusion rate was observed, with 16% of patients excluded due to a lack of an echocardiogram or severe technical problems. The authors discuss these limitations, including the possibility of selection bias.

The current study emphasizes the importance of case-specific CTP acquisition protocols and brings us one step closer to improved CTP diagnostic accuracy. Further studies are needed to optimize CTP acquisition and processing protocols, examine possible needed adjustments for other specific cases including carotid occlusion, and improve CTP resolution for the detection of small, lacunar infarcts.

REFERENCES

1. Campbell BC, Majoie CBL, Albers GW, et al; HERMES collaborators. **Penumbra imaging and functional outcome in patients with anterior circulation ischaemic stroke treated with endovascular thrombectomy versus medical therapy: a meta-analysis of individual patient-level data.** *Lancet Neurol* 2019;18:46–55 [CrossRef Medline](#)
2. Powers WJ, Rabinstein AA, Ackerson T, et al; American Heart Association Stroke Council. **2018 Guidelines for the Early Management of Patients with Acute Ischemic Stroke: A Guideline for Healthcare Professionals from the American Heart Association/American Stroke Association.** *Stroke* 2018;49:e46–110 [CrossRef Medline](#)
3. Biesbroek JM, Niesten JM, Dankbaar JW, et al. **Diagnostic accuracy of CT perfusion imaging for detecting acute ischemic stroke: a systematic review and meta-analysis.** *Cerebrovasc Dis* 2013;35:493–501 [CrossRef Medline](#)
4. Vagal A, Wintermark M, Nael K, et al. **Automated CT perfusion imaging for acute ischemic stroke: Pearls and pitfalls for real-world use.** *Neurology* 2019;93:888–98 [CrossRef Medline](#)
5. Heit JJ, Wintermark M. **Perfusion computed tomography for the evaluation of acute ischemic stroke: strengths and pitfalls.** *Stroke* 2016;47:1153–58 [CrossRef Medline](#)
6. Sotoudeh H, Bag AK, Brooks MD. **“Code-stroke” CT perfusion; challenges and pitfalls.** *Acad Radiology* 2019;26:1565–79 [CrossRef Medline](#)
7. Copen WA, Deipolyi AR, Schaefer PW, et al. **Exposing hidden truncation-related errors in acute stroke perfusion imaging.** *AJNR Am J Neuroradiol* 2015;36:638–45 [CrossRef Medline](#)
8. Borst J, Marquering HA, Beenen LF, et al; MR CLEAN investigators. **Effect of extended CT perfusion acquisition time on ischemic core and penumbra volume estimation in patients with acute ischemic stroke due to a large vessel occlusion.** *PLoS One* 2015;10:e0119409 [CrossRef Medline](#)
9. Mangla R, Ekholm S, Jahromi BS, et al. **CT perfusion in acute stroke: know the mimics, potential pitfalls, artifacts, and technical errors.** *Emerg Radiol* 2014;21:49–65 [CrossRef Medline](#)
10. Kasasbeh AS, Christensen S, Straka M, et al. **Optimal computed tomographic perfusion scan duration for assessment of acute stroke lesion volumes.** *Stroke* 2016;47:2966–71 [CrossRef Medline](#)
11. Hartman JB, Morran S, Zhu C, et al. **Use of CTA test dose to trigger a low cardiac output protocol improves acute stroke CTP data analyzed with RAPID software.** *AJNR Am J Neuroradiol* 2022 Feb 17. [Epub ahead of print] [CrossRef](#)

 J. Molad

Department of Stroke & Neurology
Tel-Aviv Sourasky Medical Center
Tel-Aviv, Israel

<http://dx.doi.org/10.3174/ajnr.A7457>