

**Online Supplemental Table: Demographic and clinical information of skull base osteomyelitis**

<i>Demographic</i>	
Sex	Male = 4, Female = 4
Median age (years [range])	69.5 (60–76)
<i>Clinical</i>	
Presence of ear or sinus lesions	
Ear lesions	4/8 (50.0%)
Sinus lesions	2/8 (25.0%)
None	2/8 (25.0%)
Diabetes mellitus	
Yes	8/8 (100.0%)
Presence of cranial nerve symptoms	
Yes	6/8 (75.0%)
No	2/8 (25.0%)
Cranial nerve symptoms	
XII	3/12 (25.0%)
X	2/12 (16.7%)
VII	2/12 (16.7%)
IX	1/12 (8.3%)
VIII	1/12 (8.3%)
VI	1/12 (8.3%)
V	1/12 (8.3%)
II	1/12 (8.3%)
Pathology	
Inflammation	4/8 (50.0%)
Chronic inflammation	2/8 (25.0%)
Granulation tissue	1/8 (12.5%)
None	1/8 (12.5%)
Cultured causative bacteria	
Pseudomonas aeruginosa	3/8 (37.5%)
Staphylococcus aureus	2/8 (25.0%)
Negative	3/8 (37.5%)
Treatment	
Antibiotics	7/8 (87.5.0%)
Antibiotics + Debridement	1/8 (12.5%)
Clinical course	
Improvement	8/8 (100.0%)