

Table 2. Comparison of collateral scores and outcome measures.

Variable	Status	n=98	Final infarct (mL), median (IQR)	Final Infarct P value	Infarct growth (mL), median (IQR)	Infarct Growth P value	Functional outcome*		Functional Outcome Odds Ratio
							mRS 0-2 (n=29)	mRS 3-6 (n=48)	
PCI	Sufficient (≥ 62)	47	19.0 (8.0–41.0)	0.01	1.0 (0.0–5.5)	0.03	21	15	2.83
	Insufficient (PCI <62)	51	37.0 (10.0–63.0)		10.0 (1.0–26.5)		8	33	
HIR	Sufficient (≤ 0.4)	44	18.0 (3.9–42.2)	0.03	3.3 (0.0–13.0)	0.21	20	27	1.55
	Insufficient (>0.4)	54	37.0 (10.5–62.5)		3.0 (0.0–25.0)		9	21	
DSA	Sufficient (ASTIN ≥ 3)	43	19.0 (5.0–42.0)	0.01	2.0 (0.0– 8.0)	0.01	13	21	0.96
	Insufficient (ASTIN <3)	55	29.0 (10.0–63.0)		6.70 (1.0– 27.2)		16	27	

* Functional outcome data (90-day mRS) only available in 77 patients