

Supplementary Table 1: Comparison of the incidence of AWE and ACE between groups of different coiled intracranial aneurysms with different follow-up intervals (months).

	≤6 months	7 to 12 months	13 to 24 months	25 to 48 months	49 to 72 months	73 to 96 months	>96 months
AWE/ Total	6/8 (75.0%)	4/7 (57.1%)	5/10 (50.0%)	4/16 (25.0%)	3/6 (50.0%)	3/6 (50.0%)	2/9 (28.6%)
ACE/ Total	5/8 (62.5%)	4/7 (57.1%)	4/10 (40.0%)	9/16 (56.3%)	4/6 (66.7%)	3/6 (50.0%)	5/9 (71.4%)

AWE: Aneurysm wall enhancement, ACE: Aneurysm central enhancement

Supplementary Table 2: Comparison of the incidence of AWE and ACE between different coiled intracranial aneurysms with and without recanalization with regards to recanalization size in millimetres (mm).

	0 mm	1 to 3 mm	4 to 5 mm	≥6 mm
AWE/ Total	6/22 (27.3%)	5/20 (25%)	10/12 (83.3%)	6/6 (100%)
p-value		0.8671	0.0033	0.0025
ACE/ Total	12/22 (54.6%)	6/19 (31.6%)	11/12 (91.7%)	5/6 (83.3%)
p-value		0.2452	0.0527	0.3547

AWE: Aneurysm wall enhancement, ACE: Aneurysm central enhancement

Supplementary Table 3: Comparison of the incidence of AWE and ACE between ruptured and unruptured coiled intracranial aneurysms.

	aSAH (n=27)	No aSAH (=33)	p-value
AWE	8 (13.3%)	19 (31.7%)	0.03
ACE	15 (25.4%)	19 (32.2%)	0.767

AWE: Aneurysm wall enhancement, ACE: Aneurysm central enhancement, aSAH: aneurysmal subarachnoid haemorrhage