

On-line Table 1: Estimated marginal means for CCA-IMT in the 2 ethnic groups^a

Adjusted for	Mean IMT (mm)		Mean Maximum IMT (mm) ^v	
	Black Individuals	White Individuals	Black Individuals	White Individuals
Age and sex	0.804 ± 0.008	0.750 ± 0.008	0.940 ± 0.020	0.770 ± 0.020
+ Cardiovascular risk factors	0.799 ± 0.008	0.755 ± 0.008	0.940 ± 0.020	0.770 ± 0.020
+ Socioeconomic status	0.800 ± 0.008	0.754 ± 0.009	0.940 ± 0.020	0.780 ± 0.020
+ CCA diameter	0.806 ± 0.008	0.749 ± 0.009	0.950 ± 0.020	0.770 ± 0.020

^a Cardiovascular risk factors: hypertension, diabetes, smoking pack-years, cholesterol, and BMI. Values are mean IMT ± SE. Differences between black and white are all $P < .001$.

On-line Table 2: Relationship between cardiovascular risk factors and mean maximum CCA-IMT in the 2 ethnic groups from univariate and multivariate regression analysis

Risk Factor	Black Individuals				White Individuals			
	Univariate		Multivariate ($R^2 = 0.11$) ^a		Univariate		Multivariate ($R^2 = 0.20$) ^a	
	B	P	B	P	B	P	B	P
Age	0.001	.000 ^b	0.001	.000 ^b	0.001	.000 ^b	0.001	.000 ^b
Male sex	0.005	.217	0.001	.768	0.014	.000 ^b	0.015	.000 ^b
Hypertension	0.011	.010 ^b	0.006	.182	0.009	.002 ^b	0.002	.592
DM	0.014	.005 ^b	0.009	.085	0.012	.052 ^b	0.013	.021 ^b
BMI	0.000	.929	0.000	.744	0.000	.792	0.000	.956
Cholesterol	0.003	.161	0.003	.068	0.001	.648	0.002	.172
Smoking: pack-years	0.000	.138	0.000	.322	0.000	.000 ^b	0.000	.054
Low social class	−0.002	.790	−0.007	.231	0.009	.006 ^b	0.005	.099

^a R^2 = proportion of variance in IMT explained by conventional risk factors.

^b $P = 0.05$.