

On-line Table: Demographic and clinical data of patients who underwent shunt surgery^a

	Subjects of the Study		Patients Who Dropped Out of the Study		P Value
	No.		No.		
Age (yr)	60	76.4 (3.8)	43	76.9 (4.6)	.569 ^b
Sex, male	60	34 (57%)	43	28 (65%)	.388 ^c
Education (yr)	60	10.2 (3.0)	43	10.6 (3.3)	.417 ^b
Duration of symptoms (yr)	60	3.3 (1.6)	43	3.3 (1.4)	.876 ^b
LP shunt	60	23 (38%)	43	9 (21%)	.060 ^c
iNPHGS total (12)	60	6.3 (1.9)	39	7.0 (1.9)	.076 ^d
iNPHGS gait (4)	60	2.3 (0.7)	39	2.3 (0.8)	.803 ^d
iNPHGS cognitive (4)	60	2.3 (0.8)	39	2.7 (0.8)	.025 ^{d,e}
iNPHGS urinary (4)	60	1.7 (1.1)	39	2.1 (1.2)	.101 ^d
TUG	55	22.1 (30.1)	33	15.8 (8.9)	.418 ^d
MMSE (30)	60	21.7 (4.6)	38	19.0 (5.3)	.011 ^{d,e}

Note:—LP indicates lumboperitoneal.

^a Data on age, education, duration of symptoms, iNPHGS, TUG, and MMSE are means (SDs). Data for the other variables are number of patients (%). Numbers in parentheses in column 1 refer to the maximum scores for the tests.

^b *t* test.

^c χ^2 test.

^d Mann-Whitney *U* test.

^e *P* value < .05.