

On-line Table: Baseline characteristics of aneurysms showing minor recanalization 6 months postembolization (N = 65)

Variables	No (%)
Sex	
Male/female	17 (26.2%)/48 (73.8%)
Age (yr)	57.9 ± 11.0
HTN	39 (60.0%)
DM	3 (4.6%)
Hyperlipidemia	6 (9.2%)
Smoking	9 (13.8%)
Antiplatelet maintenance	35 (53.8%)
Locations	
Anterior circulation	56 (86.2%)
ICA	16
AcomA and ACA	21
MCA	9
PcomA	10
Posterior circulation	9 (13.8%)
Aneurysm type	
Sidewall	15 (23.1%)
Bifurcation	50 (76.9%)
Aneurysm size	
≤7 mm	45 (69.2%)
>7 mm	20 (30.8%)
Presentation	
UIA	49 (75.4%)
SAH	16 (24.6%)
Initial occlusion	
Successful	54 (83.1%)
Residual sac	11 (16.9%)
Depth-to-neck ratio	
≤1	27 (41.5%)
>1	38 (58.5%)
Aneurysm status	
Initial	48 (73.8%)
Recanalized	17 (26.2%)
Packing density ^a	32.1 ± 9.2%
Stent	19 (29.2%)
HydroSoft coil	
≤50%	45 (69.2%)
>50%	20 (30.8%)

Note:—AcomA indicates anterior communicating artery; ACA, anterior cerebral artery; PcomA, posterior communicating artery.

^a Packing density estimated, except in 17 re-embolized aneurysms (second coiling).