

On-line Table: Diagnostic performance of the LNTA parameters in differentiating benign and metastatic lymph nodes

Protocol	Reader	AUC (95% CI)	SE of AUC	Optimal Threshold	Cross-Validated AUC
LNTA					
A	Reader 1	0.65 (0.54–0.72)	0.05	111	0.63
	Reader 2	0.63 (0.52–0.70)	0.05	113	0.61
B	Reader 1	0.73 (0.65–0.82)	0.05	99	0.72
	Reader 2	0.74 (0.64–0.80)	0.04	103	0.73
C	Reader 1	0.92 (0.85–0.96)	0.03	96	0.91
	Reader 2	0.88 (0.81–0.94)	0.04	99	0.87
Normalized LNTA to CCA					
A	Reader 1	0.71 (0.62–0.78)	0.05	0.61	0.69
	Reader 2	0.68 (0.60–0.77)	0.06	0.63	0.65
B	Reader 1	0.68 (0.59–0.77)	0.05	0.33	0.66
	Reader 2	0.76 (0.67–0.83)	0.05	0.34	0.74
C	Reader 1	0.91 (0.84–0.96)	0.03	0.31	0.90
	Reader 2	0.89 (0.82–0.95)	0.03	0.36	0.88
Normalized LNTA to IJV					
A	Reader 1	0.73 (0.64–0.81)	0.05	0.57	0.72
	Reader 2	0.66 (0.50–0.69)	0.06	0.61	0.55
B	Reader 1	0.69 (0.60–0.77)	0.05	0.35	0.68
	Reader 2	0.73 (0.63–0.80)	0.05	0.34	0.74
C	Reader 1	0.89 (0.81–0.94)	0.04	0.38	0.88
	Reader 2	0.85 (0.76–0.91)	0.04	0.43	0.84
Normalized LNTA to paraspinal muscle					
A	Reader 1	0.70 (0.60–0.78)	0.05	1.34	0.68
	Reader 2	0.63 (0.52–0.71)	0.06	1.53	0.60
B	Reader 1	0.72 (0.64–0.80)	0.05	1.55	0.71
	Reader 2	0.78 (0.70–0.86)	0.04	1.90	0.78
C	Reader 1	0.91 (0.85–0.96)	0.03	1.42	0.90
	Reader 2	0.90 (0.82–0.95)	0.03	1.55	0.88

Note:—SE indicates standard error.