

On-line Table: Numbers and percentages of various qualitative assessments among symptomatic and asymptomatic ears in hVFA and CFA according to the 2 readers

	hVFA										CFA								
	Symptomatic (n = 19)					Asymptomatic (n = 13)					Total (n = 32)		Symptomatic (n = 19)		Asymptomatic (n = 13)		Total (n = 32)		
	Reader A		Reader B		κ	Reader A		Reader B		Reader A	Reader B	Readers A and B		Readers A and B		Readers A and B	κ		
U-S distinction	11	57.9%	10	52.6%	8	61.5%	9	69.2%	19	59.4%	19	59.4%	10	52.6%	12	92.3%	22	68.8%	1
Posterior ampulla	11	57.9%	11	57.9%	9	69.2%	11	84.6%	20	62.5%	22	68.8%	19	100.0%	13	100.0%	32	100.0%	1
Lateral ampulla	9	47.4%	7	36.8%	6	46.2%	4	30.8%	15	46.9%	11	34.4%	19	100.0%	13	100.0%	32	100.0%	1
EH	10	52.6%	7	36.8%	3	23.1%	3	23.1%	13	40.6%	10	31.3%	12	37.5%	2	15.4%	14	43.8%	1
BLB impairment	6	21.1%	6	21.1%	0	15.4%	0	0.0%	6	18.8%	4	12.5%	8	42.1%	0	0.0%	8	28.1%	1

Note:—U-S distinction indicates distinction between utricle and saccule.