

On-line Table: Patient characteristics stratified by TMM versus no TMM, ICV < 70 mL versus ICV ≥ 70 mL, and ICV < 51 mL versus ICV ≥ 51 mL

| Characteristic | TMM | | ICV | | ICV | | | | |
|---------------------------------------|-------------------|---------------------|-------|-----------------------|------------------------|-------|-----------------------|----------------------|-------|
| | Positive | Negative | P | <70 mL | ≥70 mL | P | <51 mL | ≥51 mL | P |
| Patients, n | 99 | 39 | | 108 | 30 | .93 | 93 | 45 | |
| Age, (median) (IQR) (yr) | 66 (57.25, 75) | 71.5 (62.5, 77.25) | .175 | 66 (57.25, 75) | 73 (64.5, 77.25) | .083 | 68 (58, 75) | 70 (59, 76.25) | .436 |
| Males, n (%) | 63 (63.6) | 23 (59) | .611 | 66 (61.1) | 20 (66.7) | .579 | 57 (61.3) | 29 (64.4) | .72 |
| Onset-to-CT time (median) (IQR) (min) | 124.5 (64, 217) | 110 (66, 170) | .522 | 124.5 (67.25, 221.25) | 104 (58.75, 104) | .194 | 118.5 (64.75, 209.25) | 117.5 (62.5, 188.25) | .595 |
| Hypertension, n (%) | 62 (62.6) | 29 (74.4) | .19 | 68 (63) | 23 (76.7) | .161 | 56 (60.2) | 35 (77.8) | .041 |
| Diabetes, n (%) | 29 (39.3) | 16 (41) | .186 | 30 (27.8) | 15 (50) | .022 | 26 (28) | 19 (42.2) | .094 |
| Smoking, n (%) | 40 (40.4) | 13 (33.3) | .442 | 42 (38.9) | 11 (36.7) | .825 | 38 (40.9) | 15 (33.3) | .394 |
| Previous stroke and/or TIA, n (%) | 17 (17.2) | 9 (23.1) | .424 | 17 (15.7) | 9 (30) | .077 | 14 (15.1) | 12 (26.7) | .102 |
| Atrial fibrillation, n (%) | 35 (35.4) | 19 (48.7) | .147 | 40 (37) | 14 (46.7) | .339 | 34 (36.6) | 20 (44.4) | .374 |
| Dyslipidemia, n (%) | 21 (21.2) | 3 (7.7) | .059 | 21 (19.4) | 3 (10) | .227 | 20 (21.5) | 4 (8.9) | .093 |
| SBP (median) (IQR) (mm Hg) | 149 (134.25, 160) | 147.5 (120, 172.25) | .838 | 149 (132.5, 160.75) | 147.5 (119.75, 177.25) | .964 | 146.5 (130, 160) | 151.5 (126, 169.75) | .188 |
| DBP (median) (IQR) (mm Hg) | 83 (80, 91.75) | 87 (70, 92) | .442 | 83 (80, 91) | 87 (69.5, 92.25) | .637 | 83 (80, 91.75) | 86 (74, 92) | .817 |
| ABG (median) (IQR) (mol/L) | 7.2 (5.9, 9.85) | 7.6 (6.45, 8.55) | .425 | 7.15 (5.9, 9.6) | 7.75 (6.57, 9.4) | .162 | 7.1 (5.9, 9.6) | 7.7 (6.68, 9.4) | .131 |
| Baseline NIHSS score (median) (IQR) | 12.5 (9, 17) | 19 (5.75, 21) | <.001 | 13 (9, 13) | 19 (15.75, 22) | <.001 | 13 (9, 17) | 18 (13.5, 20.25) | <.001 |
| Occluded site, n (%) | | | .089 | | | .003 | | | .003 |
| MCA | 68 (68.7) | 19 (48.7) | | 76 (70.4) | 11 (36.7) | | 67 (72) | 20 (44.4) | |
| ICA | 27 (27.3) | 17 (43.6) | | 28 (25.9) | 16 (53.3) | | 24 (25.8) | 20 (44.4) | |
| MCA + ICA | 4 (4) | 3 (7.7) | | 4 (3.7) | 3 (10) | | 2 (2.2) | 5 (11) | |
| 90-day mRS 0 to ~2, n (%) | 43 (43.4) | 11 (28.3) | .099 | 48 (44.4) | 6 (21.7) | .015 | 93 (67.4) | 10 (22.2) | .005 |
| rFTD grade ≤1, n (%) | 79 (79.8) | 13 (33.3) | <.001 | 85 (78.7) | 7 (23.3) | <.001 | 75 (80.6) | 17 (37.8) | <.001 |

Note:—SBP indicates systolic blood pressure; DBP, diastolic blood pressure; ABG, admission blood glucose.