

On-line Table: Comparison of demographic data, vascular risk factor profile, clinical parameters, and collateral circulation with stroke severity^a

Parameters	NIHSS < 5 (n = 16)	NIHSS ≥ 5 (n = 49)	P Value
Age (yr) (mean)	60.7 ± 10.3	55.8 ± 11.8	.255
Male sex	16 (100)	44 (89.8)	1.000
Hypertension	8 (50)	21 (42.9)	.618
Diabetes mellitus	4 (25)	13 (26.5)	1.000
Dyslipidemia	1 (6.3)	3 (6.1)	1.000
Current smoking	10 (62.5)	30 (61.2)	.927
Prior ischemic event	3 (18.8)	11 (22.4)	1.000
Atherosclerotic ICA occlusion	12 (75)	31 (63.3)	.389
Dissection/embolic ICA occlusion	4 (25)	18 (36.7)	
Territorial (MCA) infarct	2 (12.5)	22 (44.9)	<.001
Hemodynamic infarct	2 (12.5)	21 (42.9)	
Other small infarcts	12 (75)	6 (12.2)	
ASPECTS ≥ 8	16 (100)	27 (55.1)	.001
AcomA, poor flow	7 (43.8)	24 (49)	.716
AcomA, good flow	9 (56.2)	25 (51)	
Ipsilateral PcomA, poor flow	9 (56.2)	30 (61.2)	.724
Ipsilateral PcomA, good flow	7 (43.8)	19 (38.8)	
Contralateral PcomA, poor flow	11 (68.8)	32 (65.3)	.800
Contralateral PcomA, good flow	5 (31.2)	17 (34.7)	
Ipsilateral OA, absent	3 (18.8)	23 (46.9)	.046
Ipsilateral OA, present	13 (81.2)	26 (53.1)	
Contralateral OA, absent	2 (12.5)	5 (10.2)	1.000
Contralateral OA, present	14 (87.5)	44 (89.8)	
LC, poor flow	0 (0)	18 (36.7)	<.001
LC, good flow	16 (100)	31 (63.3)	
No. of collaterals, ≥2	15 (93.7)	29 (59.2)	.010

^a Data are number and percentage unless otherwise indicated.